


# Rapid progressors in Alzheimer's Disease: Using neuroimaging to predict cognitive decline

Madison Shyer<sup>1,2</sup>, Kristofer Harris, MPH, RN<sup>2</sup>, Livia Merrill, PhD<sup>2</sup>, and Paul E. Schulz, MD<sup>1,2</sup>

(1) The University of Texas MD Anderson Cancer Center UTHHealth Houston Graduate School of Biomedical Sciences  
(2) UTHHealth Houston McGovern Medical School Dept of Neurology, Houston, TX, USA

## BACKGROUND & RATIONALE

- Alzheimer's Disease (AD) is a neurodegenerative disorder characterized by **cognitive decline**, **cortical atrophy**, and the buildup of **amyloid plaques** and **tau tangles**.
- Diffusion tensor imaging (DTI)** can be used to detect degradation of white matter in the brain as an early indicator of disease progression.
- Variability in disease progression impedes **effective identification of and treatment development for** patient subgroups who rapidly progress (**rapid progressors, RPs**).
- RPs may demonstrate **different microstructural changes** than non-RPs.




This study explores whether changes in **specific white-matter pathways** predict **cognitive score changes** between rapidly progressing (RP) and traditionally progressing (nRP) patients.

## COHORT CHARACTERISTICS

|                                                                            | Non-Rapid Progressor<br>(N = 90, 83.3%) | Rapid Progressor<br>(N = 18, 16.7 %) |
|----------------------------------------------------------------------------|-----------------------------------------|--------------------------------------|
| Age at baseline mean (± SD)                                                | 67.7 (±10.3)                            | 72.1 (±9.2)                          |
| Sex                                                                        |                                         |                                      |
| Female                                                                     | 43 (47.8%)                              | 9 (50.0%)                            |
| Male                                                                       | 47 (52.2%)                              | 9 (50.0%)                            |
| Ethnicity                                                                  |                                         |                                      |
| Hispanic or Latino                                                         | 48 (53.3%)                              | 14 (77.8%)                           |
| Not Hispanic or Latino                                                     | 42 (46.7%)                              | 4 (22.2%)                            |
| Race                                                                       |                                         |                                      |
| White                                                                      | 70 (77.8%)                              | 16 (88.9%)                           |
| Black or African American                                                  | 20 (22.2%)                              | 2 (11.1%)                            |
| <b>Baseline Conditions &amp; Other Imaging Metrics</b>                     |                                         |                                      |
| <i>Measures of neurocognitive and/or functional impairment mean (± SD)</i> |                                         |                                      |
| CDR-SB                                                                     | 2.8 (± 2.1)                             | 3.9 (± 3.4)                          |
| MMSE                                                                       | 24.5 (± 4.2)                            | 22.3 (± 5.3)                         |
| <b>PET SUVR</b>                                                            |                                         |                                      |
| Amyloid Global*                                                            | 1.01 (0.96-1.06)                        | 1.34 (1.23-1.44)                     |
| Tau Medial Temporal*                                                       | 1.12 (1.02-1.19)                        | 1.82 (1.66-1.97)                     |
| <b>vMRI (cm<sup>3</sup>), mean (± SD)</b>                                  |                                         |                                      |
| Whole brain volume                                                         | 1354.7 (± 132.1)                        | 1395.9 (± 60.4)                      |
| Ventricular volume                                                         | 26.7 (± 17.7)                           | 35.8 (± 13.7)                        |
| Left entorhinal cortex volume**                                            | 3.16 (± 0.43)                           | 2.72 (± 0.54)                        |
| Right entorhinal cortex volume***                                          | 3.31 (± 0.53)                           | 2.73 (± 0.55)                        |
| Left hippocampus volume**                                                  | 2.92 (± 0.48)                           | 3.44 (± 0.19)                        |
| Right hippocampus volume**                                                 | 2.94 (± 0.46)                           | 3.21 (± 0.11)                        |

\*p < 0.05, \*\* p < 0.01, \*\*\* p < 0.001

## RESULTS



## METHODS

### Definition

- Rapid progressors were defined as the top ~10% of individuals exhibiting the greatest decline in CDR-SB & MMSE over 18-24 months.

### Data Source

- Diagnosed AD subjects with ≥2 visit & MRI scans (n = 108) from the Health and Aging Brain Study-Health Disparities (HABS-HD) Cohort

### Analyses

- Linear mixed-effects models tested whether visit-level imaging values moderated longitudinal cognitive change between groups.
- Between-visit cognitive change was modeled as a function of ROI change, group, and their interaction.

### Metrics

- ROIs were z-scored before modeling. Covariates included age, sex, education, APOE status, and total brain volume.
- P-values were FDR corrected within each ROI set and cognitive outcome.

## DISCUSSION

- Identified two white matter measures that demonstrate **significant associations** with cognitive decline in RPs.
- Exploratory results suggest that changes in white matter microstructural integrity **may contribute** to rapid progression



The identified regions should be interpreted as potential candidates for further study

## FUTURE DIRECTIONS

- Investigate the relationship between **neuroanatomical changes** and **cognitive domain decline** in nRPs v. RPs
- Develop prediction models to identify potential factors (e.g., more **malignant forms of tau**) that underly rapid progression
- Inform randomized clinical trial design, patient stratification, and targeted drug development specific to RPs

Research reported on this publication was supported by the National Institute on Aging of the National Institutes of Health under Award Numbers R01AG054073, R01AG058533, R01AG070862, P41EB015922 and U19AG078109.