

Divergent scalp-to-region distance alteration patterns in autism spectrum disorders, Parkinson’s disease and Alzheimer’s disease

Liqin Yang^{1,2,#,*}, Junhao Zhang^{3,#}, Junlong Wang¹, Hsin-Hsiung Huang⁴, Hongbin Han⁵, Daniel Razansky³, for the ADNI, Axel Rominger¹, Jie Lu⁶, Ruiqing Ni^{1, 3,*}
¹Department of Nuclear Medicine, Inselspital, Bern University Hospital, University of Bern, 3010 Bern, Switzerland; ²Department of Radiology, Huashan Hospital, Fudan University, 200040 Shanghai, China; ³Institute for Biomedical Engineering, ETH Zurich & University of Zurich, 8093 Zurich, Switzerland; ⁴Department of Statistics and Data Science, University of Central Florida, Orlando, FL, USA; ⁵Department of Radiology, Peking University Third Hospital, 10091 Beijing, China; ⁶Department of Radiology, Xuanwu Hospital, Beijing, China

- An automatic pipeline enables scalp-to-region distance (SRD) measurement and facilitates brain stimulation planning.
- ASD patients show widespread SRD decreases, especially in the orbitofrontal cortex and adjacent regions.
- AD patients present increased SRD in the frontal gyrus and decreased SRD in the parahippocampal gyrus.

INTRODUCTION


Brain stimulation is increasingly recognized as an effective therapeutic intervention for many brain diseases. Distance between the scalp and brain regions is pivotal for neurostimulation planning, but its alterations in brain diseases are largely unknown.

ABIDE	CN	ASD	p-value
N	381	313	
Sex (M/F)	337/44	258/55	0.024
Age (years)	15.88±6.34	15.7±5.69	0.6883
PPMI	CN	PD	p-value
N	84	225	
Sex (M/F)	55/29	144/81	0.8095
Age (years)	59.5±11.62	61.73±9.49	0.085
ADNI	CN	AD	p-value
N	224	155	
Sex (M/F)	114/110	75/80	0.6315
Age (years)	76.04±6.42	74.66±7.43	0.0534

We developed an automatic pipeline to calculate scalp-to-region distance (SRD) values from T1 MR images and applied

METHODS


it to 1382 participants, including patients with autism spectrum disorder (ASD), Parkinson’s disease (PD), Alzheimer’s disease (AD), and cognitively normal controls (CNs).




Region Label	SRD value
R_superior_frontal_gyrus	27.99
L_superior_frontal_gyrus	27.62
R_superior_frontal_gyrus	20.95
L_superior_frontal_gyrus	19.22

RESULTS

Compared with CNs, ASD patients showed widespread SRD decreases with prominent involvement of the frontal lobe, especially the orbitofrontal cortex and adjacent regions.



In contrast, in AD patients, significantly increased SRD values were observed in the frontal gyrus. No significant SRD alteration was found in PD patients.



CONCLUSIONS

The automatic SRD calculation pipeline and the different patterns of SRD alterations in these diseases may support future validation studies for neurostimulation planning.

Contact:
ruiqing.ni@unibe.ch

REFERENCES

Zhang J, et al. Brain Stimul. 2023 Mar-Apr;16(2):653-656. doi: 10.1016/j.brs.2023.03.011.

Yang L, et al. bioRxiv [Preprint]. 2026 May 18:2026.05.14.725296. doi:10.64898/2026.05.14.725296.

ALZHEIMER'S ASSOCIATION | AAIC 26