

Long-term longitudinal patterns of hypometabolism and atrophy progression from preclinical to dementia stages of Alzheimer's disease

Miguel Labrador-Espinosa¹, Jesús Silva-Rodríguez¹, Michael Schöll²*, Michel Grothe¹*

1. Reina Sofia Alzheimer Center, CIEN Foundation, ISCIII, Madrid, Spain. 2. University of Gothenburg, Department of Psychiatry and Neurochemistry, Mölndal, Sweden.

Poster IDs

AAIC: #4694; AIC: #12338

INTRODUCTION

- Structural MRI-derived atrophy and FDG-PET-measured hypometabolism are complementary markers of neurodegeneration in Alzheimer's disease (AD), capturing cumulative tissue loss and synaptic-neuronal dysfunction, respectively.
- However, the temporo-spatial evolution of cortical hypometabolism and atrophy patterns from preclinical stages to AD dementia remains incompletely understood.
- Aim:** In this study, we investigated the temporal evolution of cortical hypometabolism and atrophy using serial FDG-PET and MRI acquisitions in individuals who developed AD dementia.

METHODS

- Participants:** 208 nondemented individuals (15% CU, 85% MCI) enrolled in the ADNI study who developed biomarker-confirmed AD dementia over clinical follow-up (Table 1).
 - All participants underwent serial MRI and [18F]FDG-PET acquisitions spanning from -15 years before to +5 years after dementia onset.
- Neuroimaging:**
 - Cortical thickness on MRI and standardized uptake value ratios on FDG-PET were calculated across 200 cortical regions defined in the Schaefer atlas.
 - Region-wise z-scores were calculated to quantify cortical thickness and hypometabolism relative to healthy elderly controls (N(MRI)=218, N(FDG)=56).
- Statistical analyses:**
 - Longitudinal atrophy and hypometabolism progression of each brain region was calculated using linear mixed-effects models adjusted for age and sex.
 - Predicted z-scores were generated at 2-year intervals spanning from -10 years before dementia onset to +2 years after.
 - Inter-modality z-score differences quantified the relative predominance of regional hypometabolism versus atrophy over the course of disease progression.




Figure shows predicted z-scores for hypometabolism on FDG-PET (a), cortical thinning on MRI (b) and inter-modality z-score differences (c) at 2-year intervals across a 12-year window spanning 10 years before dementia onset to 2 years after.

- Hypometabolism on FDG-PET**
 - ✓ First pronounced hypometabolism ($z > 1$) was observed in the posterior cingulate cortex and precuneus approximately 3.5 years before dementia, temporally overlapping with MCI onset.
 - ✓ Over progression to dementia, hypometabolism extended to lateral temporo-parietal areas.
- Cortical atrophy on MRI**
 - ✓ By contrast, first pronounced cortical thinning was observed in the temporal pole and inferior temporal cortex at a comparable time point, approximately 3.5 years before dementia.
 - ✓ Cortical thinning progressively involved additional fronto-temporo-parietal regions during the progression to dementia.
- Divergent regional patterns**
 - ✓ A marked predominance of cortical hypometabolism over atrophy was already evident in a widespread cortical pattern at the preclinical stage (-9.3 ± 3.1 years).
 - ✓ As individuals progressed to MCI and dementia, a divergent regional profile emerged, characterized by prominent hypometabolism in medial parietal regions and predominant atrophy in inferior temporal regions.

CONCLUSION

- Cortical atrophy and hypometabolism in AD progression show markedly different origins and temporal trajectories, indicating differential patterns of neurodegenerative phenomena over the course of AD.

REFERENCES

[1] Jack Jr. CR, Andrews JS, Beach TG, et al. Revised criteria for diagnosis and staging of Alzheimer's disease: Alzheimer's Association Workgroup. *Alzheimer's & Dementia*. 2024;20(8):5143-5169. doi:10.1002/alz.13859.

[2] Kljajevic V, Grothe MJ, Ewers M, Teipel S; Alzheimer's Disease Neuroimaging Initiative. Distinct pattern of hypometabolism and atrophy in preclinical and predementia Alzheimer's disease. *Neurobiol Aging*. 2014;35(9):1973-1981. doi:10.1016/j.neurobiolaging.2014.04.006

[3] Villain N, Fouquet M, Baron JC, et al. Sequential relationships between grey matter and white matter atrophy and brain metabolic abnormalities in early Alzheimer's disease. *Brain*. 2010;133(11):3301-3314. doi:10.1093/brain/awq203.

CONTACT



Miguel A. Labrador-Espinosa
mlabrador@fundacioncien.es
X (Twitter): @mlabesp

The statistical values are mean (SD) for numerical variables, count (%) for categorical variables, and median [interquartile range: p25, p75] for nominal variables. Abbreviations: CU = Cognitively unimpaired; MCI = Mild cognitively impaired.