


BACKGROUND


Recent research has suggested that Neuropsychiatric Symptoms (NPS) may be related to AD pathogenesis across the cognitive impairment continuum. However, the role of plasma biomarkers regarding NPS in preclinical stages is not clear. We examined the relationships between NPS and p-tau181 and Aβ42 in a sample of Cognitively Unimpaired (CU) and Subjective Cognitive Decline (SCD) participants.



METHODS

A total of 296 participants aged +50 years from the Compostela Ageing Study (CompAS) recruited from primary care health centers were diagnosed as CU (N=117), and SCD (N=179) according to [Jessen et al. \(2014\)](#), and defined by cognitive complaints exceeding age-adjusted cut-off scores in the Questionnaire d’Autoévaluation de la Mémoire (QAM) (5<sup>th</sup> percentile, [Pereiro et al., 2021](#)). NPS were assessed using the Mild Behavioral Impairment-Checklist (MBI-C) 12-item (proposed by [Mallo et al., 2023](#)).

Figure 1. Figure 2. Coordinates for the 12 MBI-C items and quadrants proposed by [Mallo et al. \(2023\)](#).



Legend: Q-I: Quadrant I; Q-II: Quadrant-II; Q-III: Quadrant-III; Q-IV: Quadrant-IV, MBI-C: Mild Behavioral Impairment- Checklist.

RESULTS

Plasma biomarkers, p-tau181 and Aβ42, were obtained. Aβ42 concentrations were measured using the Simoa Human Neurology 3-Plex A Advantage Kit, while plasma levels of p-tau 181 were determined using the ALZpath Simoa® p-tau 217 v2 Assay. Group differences were examined using ANOVA with age, years of schooling, MBI-C 12-item total score and each MBI-C quadrant score as dependent variables, and clinical diagnosis as factor. Linear regression analyses (LRA) were performed with MBI-C 12-item total score and scores in each quadrant, Quadrant-I (Q-I, Resistance), Quadrant-II (Q-II, Restlessness), Quadrant-III (Q-III Desolation), Quadrant-IV (Q-IV Fattening), as dependent variables; p-tau181 and Aβ42 were predictors.

CONCLUSIONS

P-tau181 and Aβ42 were associated with NPS (total score, Q-I and Q-II) across all participants. However, group-specific patterns emerged: p-tau181 was linked to these NPS in SCD, while Aβ42 was associated in CU.

TAKE-HOME MESSAGE

Tau and amyloid may differentially contribute to NPS depending on cognitive status, particularly those related to Resistance and Restlessness domains.

REFERENCES

Jessen, F., et al. (2014). A conceptual framework for subjective cognitive decline in preclinical Alzheimer’s disease. *Alzheimer’s & Dementia*, 10(6), 844–852. <https://doi.org/10.1016/j.jalz.2014.01.001>

Mallo, S. C. et al. (2023). Assessment of Mild Behavioral Impairment: Methodological issues. *International Psychogeriatrics*, 35(S1), 25–26. <https://doi.org/10.1017/S1041610223001874>

Pereiro, A. X., et al. (2021). Complaint severity predicts progression from SCD to MCI: A machine learning approach. *Journal of Alzheimer’s Disease*, 82(3), 1229–1242. <https://doi.org/10.3233/JAD-210334>

Table 1. One-way ANOVA comparing age, education, and MBI-C 12-item total and MBI-C quadrant scores across clinical diagnostic groups CU and SCD. Means and SD (in parentheses).

| Variable              | CU (N=117)   | SCD (N=179)  | F(1,294) | p    |
|-----------------------|--------------|--------------|----------|------|
| Age                   | 63.53 (8.82) | 66.48 (8.80) | 7.91     | .005 |
| Years of schooling    | 14.00 (5.59) | 12.96 (5.62) | 2.41     | .121 |
| MMSE                  | 28.65 (1.50) | 28.33 (1.61) | 3.08     | .080 |
| MBI-C 12 items. Total | 2.57 (4.58)  | 3.59 (4.95)  | 3.20     | .074 |
| Aβ42                  | 10.49(3.91)  | 9.69(5.32)   | 1.82     | .178 |
| p-tau181              | 23.86(29.15) | 21.99(15,40) | .496     | .482 |

Legend: Q-I: Quadrant I; Q-II: Quadrant-II; Q-III: Quadrant-III; Q-IV: Quadrant-IV, MBI-C: Mild Behavioral Impairment- Checklist; MMSE: Mini Mental State Examination.

Table 2. Linear regression analyses of MBI-C total and quadrant scores with p-tau181 and Aβ42 as predictors with the entire sample (N=296).

|                      | Aβ42 |     |      |       |       | p-tau181 |     |     |      |      |
|----------------------|------|-----|------|-------|-------|----------|-----|-----|------|------|
|                      | B    | SE  | β    | t     | p     | B        | SE  | β   | t    | p    |
| MBI-C 12 items total | -.12 | .60 | -.12 | -2.03 | <.001 | .03      | .01 | .14 | 2.44 | .015 |
| Q-I Resistance       | -.03 | .01 | -.12 | -2.14 | .033  | .01      | .01 | .18 | 3.09 | .002 |
| Q-II Restlessness    | -.03 | .01 | -.16 | -2.80 | .005  | .01      | .01 | .17 | 3.06 | .002 |
| Q-III Desolation     | -.03 | .02 | -.07 | -1.28 | .202  | .01      | .01 | .10 | 1.80 | .072 |
| Q-IV Flattening      | -.01 | .01 | -.05 | -.96  | .333  | .01      | .01 | .02 | .40  | .68  |

Legend: Q-I: Quadrant I; Q-II: Quadrant-II; Q-III: Quadrant-III; Q-IV: Quadrant-IV, MBI-C: Mild Behavioral Impairment- Checklist.

Table 3. Linear regression analyses of MBI-C total and quadrant scores with p-tau181 and Aβ42 as predictors with SCD and CU participants.

|                      | Aβ42 |     |      |       |      | p-tau181 |     |     |      |       |
|----------------------|------|-----|------|-------|------|----------|-----|-----|------|-------|
|                      | B    | SE  | β    | t     | p    | B        | SE  | β   | t    | p     |
| SCD (N=179)          |      |     |      |       |      |          |     |     |      |       |
| MBI-C 12 items total | -.08 | .07 | -.09 | -1.22 | .223 | .05      | .02 | .15 | 2.06 | .040  |
| Q-I Resistance       | -.01 | .01 | -.05 | .74   | .458 | .02      | .01 | .25 | 3.43 | <.001 |
| Q-II Restlessness    | -.02 | .01 | -.10 | -1.33 | .185 | .01      | .01 | .23 | 3.06 | .003  |
| Q-III Desolation     | -.03 | .03 | -.07 | -1.01 | .310 | .01      | .01 | .07 | .99  | .319  |
| Q-IV Flattening      | -.01 | .01 | -.07 | -1.00 | .318 | .00      | .00 | .01 | .07  | .938  |
| CU (N=117)           |      |     |      |       |      |          |     |     |      |       |
| MBI-C 12 items total | -.16 | .11 | -.13 | -1.47 | .145 | .02      | .01 | .16 | 1.70 | .091  |
| Q-I Resistance       | -.08 | .03 | -.22 | -2.40 | .018 | .01      | .00 | .14 | 1.52 | .131  |
| Q-II Restlessness    | -.06 | .02 | -.25 | -2.79 | .006 | .01      | .01 | .17 | 1.94 | .055  |
| Q-III Desolation     | -.01 | .04 | -.03 | -.41  | .680 | .01      | .01 | .16 | 1.73 | .086  |
| Q-IV Flattening      | -.01 | .03 | -.01 | -.01  | .986 | .00      | .00 | .04 | .50  | .616  |

Legend: Q-I: Quadrant I; Q-II: Quadrant-II; Q-III: Quadrant-III; Q-IV: Quadrant-IV, MBI-C: Mild Behavioral Impairment- Checklist.

AAIC >26

ALZHEIMER'S ASSOCIATION

ISTAART

ALZHEIMER'S ASSOCIATION

Alzheimer’s Association International Conference | July 12 – 15, 2026 | London, UK

