




Penn Frontotemporal
Degeneration Center

Adjusting plasma p-tau₂₁₇ levels for kidney function to improve association with amyloid and tau pathology

Katheryn A.Q. Cousins, PhD¹, Christopher Brown, MD, PhD¹, Leslie M. Shaw, PhD², Magdalena Korecka, PhD², David A. Wolk, MD¹

¹Department of Neurology, ²Department of Pathology and Laboratory Medicine, Perelman School of Medicine, University of Pennsylvania; 3600 Spruce Street, Philadelphia, PA 19104.



Alzheimer's Disease
Research Center

Background

- Plasma phosphorylated tau 217 (**p-tau₂₁₇**) measured by Fujirebio Lumipulse (RUO) shows excellent performance to detect abnormal β -amyloid (**A β**), indicative of Alzheimer's disease (**AD**)
- Comorbidities may confound AD blood biomarker levels, thereby affecting performance:
 - Kidney function alters blood p-tau₂₁₇ levels independently of AD pathology: chronic kidney disease (**CKD**) (*i.e.*, low estimated glomerular filtration rate [**eGFR**]) can lead to false positives by p-tau₂₁₇
 - Conversely, higher body mass index (**BMI**) is associated with lower blood p-tau₂₁₇
- It is unclear whether strategies accounting for these conditions improve blood biomarker accuracy and interpretation
- Here we evaluate whether linear adjustments can improve blood biomarker associations with AD pathology
- Study Goals:**
 - In A β PET negative (A β -) cases, we model plasma p-tau₂₁₇ adjusted for eGFR and BMI
 - Compute adjusted p-tau₂₁₇ values in all participants
 - Test if adjusted p-tau₂₁₇ strengthens correlations with A β and tau PET, compared to unadjusted (raw) p-tau₂₁₇
 - Test if p-tau₂₁₇ adjustment improves biomarker classification

Methods & Participants

	ADNI	ADNI A β -	ADNI A β +	p
n	17	702	448	
A β PET Centiloid	--	2.5 [-4.0, 8.0]	74.0 [46.0, 102.2]	<0.001
Global TauMax	23.6 [0.2, 39.2]	0.0 [0.0, 0.0]	0.5 [0.0, 9.1]	<0.001
Tau PET SUVR	1.3 [1.2, 2.0]	1.2 [1.1, 1.2]	1.3 [1.2, 1.6]	<0.001
Cognitive Status = MCI (%)	12 (70.6%)	225 (32.1%)	279 (62.3%)	<0.001
Age at Plasma (years)	70.8 [66.0, 73.7]	71.6 [65.6, 77.0]	75.6 [70.3, 80.5]	<0.001
eGFR	79.2 [67.4, 90.6]	77.9 [65.3, 89.5]	73.3 [61.9, 85.0]	<0.001
Body Mass Index (BMI)	26.2 [23.5, 31.0]	27.1 [24.0, 30.8]	26.1 [23.4, 29.6]	0.024
Years from plasma to eGFR	0.2 [0.1, 0.3]	0.1 [0.0, 0.2]	0.1 [0.0, 0.2]	0.109
Sex = Male (%)	7 (41.2%)	302 (43.0%)	195 (43.5%)	0.972
Self-reported Race (%)				<0.001
White	9 (52.9%)	457 (66.2%)	354 (79.2%)	
Black	7 (41.2%)	148 (21.4%)	69 (15.4%)	
Other/Unknown	1 (5.9%)	85 (12.3%)	24 (5.4%)	
Ethnicity = Hispanic (%)	0	89 (12.7%)	28 (6.3%)	0.001
APOE ϵ 4 allele(s) (%)	10 (62.5%)	154 (23.4%)	286 (66.8%)	<0.001

Table 1. ADNI sample characteristics. Median [interquartile range] and Kruskal Wallis p-values are reported for continuous variables; count (Percentage) and chi-square p-values for categorical.

- Participants were selected from the Alzheimer's Disease Neuroimaging Initiative (**ADNI**) (n= 1,167)
 - Inclusion criteria: measurements for plasma p-tau₂₁₇, **eGFR calculated from creatinine**, body mass index (**BMI**), and either A β or tau PET
 - Exclude eGFR measurements >2years from plasma
 - Cognition was normal (n=651) or mild cognitive impairment (**MCI**; n=516)
 - Missing data: A β PET (n=17), Tau PET SUVR (n=221), TauMax (n=506), Race (n=13), Ethnicity (n=4), APOE ϵ 4 (n=67)
- A β PET Status: Centiloids ≥ 20 = A β +, Centiloids <20 = A β -
- Fujirebio Lumipulse measured plasma p-tau₂₁₇ at Penn
- Adjusted p-tau₂₁₇ based on model in A β -: plasma p-tau₂₁₇ (log) ~ eGFR + BMI (Figure 1)

Results: Computing eGFR+BMI-adjusted plasma p-tau₂₁₇




Figure 1. Plasma p-tau₂₁₇ associations with eGFR and BMI. (A.) Scatterplots and least squares regression lines of plasma p-tau₂₁₇ log-transformed by BMI and eGFR. Color indicates A β PET status. (B.) Linear model of p-tau₂₁₇(log) as a function of BMI and eGFR in PET A β - cases. This model was applied to the full sample to calculate adjusted values. (C.) Linear model of p-tau₂₁₇(log) as a function of BMI, eGFR, and A β and tau pathologies in all cases.

- Computed adjusted values in all participants based on model (Figure 1B)

Results: Correlations with PET A β and tau




Figure 2. Scatterplots comparing Spearman correlations with AD pathology. (A.) A β PET Centiloids correlations with plasma p-tau₂₁₇ log-transformed adjusted ([adj] green) vs. raw (yellow). (B.) Tau PET TauMax and meta temporal SUVR score for Flortaucipir correlations with plasma p-tau₂₁₇ adjusting (adj; green) vs. raw (yellow). Spearman's rho and p-value are reported.

Results: AD classification using eGFR+BMI-adjusted plasma p-tau₂₁₇




Figure 3. Changes in biomarker status, stratified by PET A β status. Panels stratify by PET Status, color indicates pairwise increase (red = Intermediate to A β +, coral = A β - to Intermediate) or decrease (light blue = A β to Intermediate, dark blue = Intermediate to A β -) of p-tau₂₁₇ adjusted (adj) compared to raw. Lines indicate 95% specificity (solid) and 95% sensitivity cutpoints (dotted) for raw (grey) and adjusted (black) p-tau₂₁₇. Annotations indicate number and mean A β Centiloid, eGFR, BMI, and age for each group.

Conclusions & Future Directions

- eGFR and BMI were significantly associated with plasma p-tau₂₁₇ concentrations, even after for controlling for age, A β and tau pathologies (Figure 1B and 1C)
- Linear adjustments of plasma p-tau₂₁₇ showed incremental improvement compared to raw values:
 - Spearman correlations with A β PET (Figure 2A) and TauMax (Figure 2B) were slightly higher for adjusted compared to raw plasma p-tau₂₁₇
 - Receiver operating curve (ROC) area under the curve was slightly higher for adjusted (AUC=0.91) than raw p-tau₂₁₇ (AUC=0.90)
 - Overall, 16 of 1150 cases had improved classification using adjustment: 1 A β case and 15 A β - cases (Figure 3)
- Future work should evaluate in populations with more severe morbidities, and test alternative strategies such as eGFR calculated by Cystatin C or non-linear adjustment

Funding

Funding from the National Institute of Aging (P01-AG066597, P30-AG072979, R01-AG087258), and the Penn Institute on Aging.

Data collection and sharing for the Alzheimer's Disease Neuroimaging Initiative (ADNI) is funded by the National Institute on Aging (National Institutes of Health Grant U19AG024904)