

Circadian dysregulation links social isolation to neurodegenerative risk: A translational study in *Drosophila melanogaster*

Javiera Palma 1,2 Christine Gierke 1,2,3,4 Martina Morales 1,2, Millaray Pizarro 1,2, Rocio Diaz 1,2,6 Vicente Iturriaga 1,2, Alfonso

Valderrama 1,2, 5 Fernanda Cabrera 1,2,5 , Walthers Diaz-Gierke1,2,7 Carlos F. Navarro 1,2 & Francisco Del Pino 1,2,8.

1 Social Isolation Research Center (SIRC), Santiago, Chile, 2. ISOFLY Lab NiBG, Biomedical Sciences Institute, University of Chile, Santiago, Chile, 3. Neurology and Neurosurgery Service, Clinical Hospital University of Chile (HCUCH), 4. School of Psychology, Diego Portales University (UDP), Santiago, Chile. 5. School of Psychology, Silva Henríquez University, 6. Pontifical Catholic University (PUC), Santiago, Chile. 7. School of Psychology, Adolfo Ibañez University, Santiago, Chile. 8. Biomedical Sciences Institute (ICBM), NiBG , Faculty of Medicine University of Chile, Santiago, Chile.

Introduction

The Clinical Paradigm: Social isolation is recognized by the Lancet Commission as a modifiable risk factor for cognitive decline and dementia, accounting for roughly 4% of global cases. Clinically, across neurodegenerative diseases, circadian disruption frequently manifests as fragmented sleep, altered rest-activity rythms, nocturnal agitaptn, and impaired behavioral timing.

Social Isolation as a Circadian Stressor: In *Drosophila melanogaster*, chronic social isolation has been shown to reduce sleep and alter sleep architecture, particularly through shorter and more fragmented rest bouts. These behavioral changes suggest that the absence of social context can destabilize sleep–wake organization.

Neurobiological Rationale: In *Drosophila melanogaster*, social isolation has been shown to reduce and fragment sleep, suggesting that disrupted rest is not only a behavioral outcome but also a potential biological stressor. Brown et al. (2017) demonstrated that isolation-induced sleep loss activates cellular stress pathways, particularly endoplasmic reticulum stress and the unfolded protein response (UPR), reflected by increased BiP/GRP78 expression. Together with evidence linking circadian disruption to oxidative imbalance and neuronal vulnerability, this supports a translational pathway connecting social isolation, sleep fragmentation, cellular stress, and neurodegenerative risk.

Methods

