



INTRODUCTION

- The prodromal phase of AD begins decades before cognitive symptoms, characterized by metabolic dysregulation, immune-metabolic vulnerability, and early amyloid accumulation^{1,2}. Targeting these systems through FDA-approved therapeutics during this window could reduce AD risk³.
- Our previous epidemiological analyses demonstrated that lipid-lowering⁴, glucose-regulating⁵, and anti-inflammatory⁶ therapeutics were each associated with lower AD risk, with the greatest benefit when used in combination⁷.
- Triple combination therapy (Triple-Rx) consisting of atorvastatin (lipid metabolism), tofacitinib (JAK inhibition), and pioglitazone (PPARγ agonism/glucose regulation), targets complementary multisystem pathways implicated in early AD pathogenesis^{8,9,10}.
- This study demonstrated that Triple-Rx attenuates early amyloid pathology (6E10) while preserving microglial homeostasis (Iba1), without inducing peripheral immune activation, in humanized (h) *APP*/*hAPOE3* and *hAPP*/*hAPOE4* mice, a model recapitulating sex- and APOE-dependent AD risk vulnerabilities^{11,12}.

Alzheimer's Disease Network




Preclinical Validation




METHODS


- 15-month female and male hAPP/hAPOE3 and hAPP/hAPOE4 mice were treated for 90 days with control chow, monotherapies (atorvastatin [ATV], tofacitinib [TOF], or pioglitazone [PGZ]), or combinatory chow regimens, including a triple-Rx: ATV+TOF+PGZ (300/250/200 mg/kg).
- Microglial-related neuroinflammatory status was assessed by Iba1 immunolabeling, while peripheral immune activation was evaluated via flow cytometry.
- Amyloid-associated pathology was assessed using 6E10 immunolabeling, and plasma A β 42:40 ratios were measured via MSD assays to assess peripheral amyloid.
- Cognitive function was assessed via novel object recognition (NOR) and body composition was assessed via EchoMRI before and after treatment.




Triple-Rx reduced amyloid-beta (A β) and microglial expression in the brain.




Triple-Rx prevents decline in cognition.



Triple-Rx prevents age-associated weight gain.



Triple-Rx increases plasma A β 42:40 in females, consistent with decline in brain while not affecting peripheral immune profiles across sexes.




RESULTS & DISCUSSION

- Iba1 immunoreactivity under triple-Rx was differentially reduced across sex, genotype, and region, demonstrating preserved microglial homeostasis without triggering neuroinflammation or peripheral immune activation.
- Triple-Rx significantly reduced 6E10 across multiple brain regions in both sexes and APOE genotypes, indicating broad attenuation of early amyloid-associated pathology in a pre-symptomatic disease context.
- Elevated plasma Aβ_{42:40} ratios in triple-Rx animals suggest enhanced peripheral amyloid clearance, alongside preserved cognitive performance and body weight trajectories, supporting the safety and tolerability of this multi-pathway strategy.
- These findings support combination therapy as a precision prevention strategy for AD and highlight IHC-based amyloid and microglial endpoints as indicators of therapeutic efficacy during early, pre-pathological disease stages.



Center for Innovation in Brain Science



REFERENCES

1. Caldwell CG, Yao J, Britton RD. Targeting the Prodromal Stage of Alzheimer's Disease: Bioenergetic and Mitochondrial Opportunities. *Neurotherapeutics*. 2015;12(1):66–80.
2. Livingston G, Huntley J, Sommerlad A, et al. Dementia prevention, intervention, and care: 2020 report of the Lancet Commission. *The Lancet*. 2020;396(10248):413–46.
3. Vitali F, Branigan GL, Britton RD. Preventing Alzheimer's disease within reach by 2025: Targeted-risk-AD-prevention (TRAP) strategy. *Alzheimers Dement Transl Res Clin Interv*. 2021;7(1):21190.
4. Torralde-Haro G, Branigan GL, Vitali F, Gellman N, Zissimopoulos JM, Britton RD. Statin therapy and risk of Alzheimer's and age-related neurodegenerative diseases. *Alzheimers Dement*. 2020;6(1):e12108.
5. Torralde-Haro G, Branigan GL, Britton RD, Rodgers KE. Association between Specific Type 2 Diabetes Therapies and Risk of Alzheimer's Disease and Related Dementias in Propensity-Score Matched Type 2 Diabetic Patients. *Front Aging Neurosci*. 2022;14:87304.
6. Cortes-Flores H, Torralde-Haro G, Britton RD. Anti-inflammatory drugs as risk modifiers for Alzheimer's disease: a step towards precision prevention. *Alzheimers Dement*. 2022;18(5):e06694.
7. Vitali F, Torralde-Haro G, Branigan GL, Vitali F, Britton RD. Combination therapy targeting Alzheimer's disease risk factors is associated with a significant delay in Alzheimer's disease-related cognitive decline. *Alzheimers Dement Transl Res Clin Interv*. 2025;11(1):e070074.
8. Yang LG, March ZM, Stephenson RA, Narayan PS, Apoliprotein E in lipid metabolism and neurodegenerative disease. *Trends Endocrinol Metab*. 2023;34(8):430–45.
9. Faqueti ML, Slappendel L, Bignon H, et al. Baricitinib and tofacitinib off-target profile, with a focus on Alzheimer's disease. *Alzheimers Dement Transl Res Clin Interv*. 2024;10(1):e12445.
10. Alhawal A, Alskhan R, Alsaid M, et al. Protective Effects of Plogizalone on Cognitive Impairment and the Underlying Mechanisms: A Review of Literature. *Drug Des Dev Ther*. 2022;16:2919–31.
11. McLean JW, Bhattrai A, Vitali F, Rakes AC, Wiegand JPL, Britton RD. Contributions of sex and genotype to exploratory behavioral differences in an aged humanized APOE mouse model of late-onset Alzheimer's disease. *Learn Mem*. 2022;29(9):321–31.
12. Qi G, Mi Y, Shi X, Gu H, Britton RD, Yin F. APOE4 impairs neuron-astrocyte coupling of fatty acid metabolism. *Alzheimers Dement*. 2020;16(S2):e045251.

ACKNOWLEDGEMENTS

Research was supported by National Institute on Aging grants T32AG06187 and P01AG026572 to RDB, and the Center for Innovation in Brain Science to RDB.
We would like to thank all the staff, members and students at the Center for Innovation in Brain Science.