


BACKGROUND

- National quality improvement initiatives have attempted to mitigate opioid-related harms in postamputation patients.
- However, the prevalence of high-risk opioid prescribing in this population and the risk factors associated with persistent opioid use after major lower limb amputation are not well described.
- Filling this knowledge gap will guide safe-opioid prescribing practices for amputation patients.

OBJECTIVE



To identify risk factors associated with prolonged opioid use after major lower limb amputation.

METHODS and DEFINITIONS

Data Source: 20% National Sample of Medicare Fee-for-service Claim data, 2008 – 2018.

Study Sample: Patients who underwent major lower limb amputation (including hip, below hip, knee, below knee, ankle, and below ankle amputation) between 1/1/2009 and 6/30/2018 identified using ICD-9, ICD-10, and CPT codes.

Exclusion Criteria: Patients that did not discharge home, or had a length of stay > 30 days, or without parts A, B, and D continuous enrollment 1 year before admission and 6 months after discharge, or had subsequent surgery based on anesthesia CPT codes for surgery 180 days after discharge.

Persistent Opioid Use: 1 opioid prescription filled in the 4-90 days after discharge AND another fill 91 – 180 days after discharge.

High-risk opioid prescribing: Overlapping opioid prescriptions, concurrent benzodiazepine and opioid prescriptions, daily-doses>100 OMEs, multiple prescribers, and new long-acting opioid-use 90-days postoperatively.

Chronic Opioid User: 180-day supply over one year and at least 1 prescription within 60 days of surgery.

Opioid Naïve User: Did not fill an opioid prescription between 365 and 31 days before admission.

Intermittent Opioid User: Filled some opioid prescriptions (<180-day supply) between 365 and 31 days before admission.




Patient Covariates: Age, sex, race, Charlson Comorbidity index, presence of mental health disorders, co-existing pain conditions, level of amputation, filled benzos, sedatives, anxiolytics within 30 days before admission, filled gabapentinoids within 30 days before admission.

Statistical Methods: We used the Cochran-Armitage-Trend-Test to determine trends in high-risk prescribing over time, and multilevel mixed-effects logistic regression to determine predictors of prolonged use.

RESULTS

Risk-factors associated with prolonged opioid use in chronic and non-chronic opioid users

Cohort	Variable	Odds-Ratio	P-value
Chronic User	Opioid filled 30 days prior to admission	2.21	0.033
	Opioid filled 30 days prior to admission	4.46	<0.001
	Age < 55 years	1.41	0.04
Non-Chronic User	Concurrent Gabapentinoid	1.92	<0.001
	Concurrent Sedative/Anxiolytic	1.75	<0.001



CONCLUSIONS

- Despite increased national awareness and attempts to improve opioid prescribing patterns in the past decade, high-risk prescribing practices remain common and have not changed over time for patients undergoing lower extremity amputation surgery.
- Following lower extremity amputation, a focus on preoperative risk factors, particularly preoperative opioid exposure, will be of high-value to optimize postamputation pain control and promote safe opioid stewardship.

CORRESPONDING AUTHOR: Theodore A. Kung, MD, Associate Professor Section of Plastic Surgery, Department of Surgery University of Michigan 1500 E. Medical Center Drive 2130 Taubman Center Ann Arbor, MI 48109-5231 Office: 734.647.3842 Email: thekung@med.umich.edu