

Background

- Ventral hernia repair in oncologic patients is challenging due to poor tissue quality, immunosuppression, malnutrition, and prior operations.
- Synthetic and biologic meshes offer advantages and disadvantages, including strength and resistance to contamination
- Biosynthetic mesh material, such as that composed of poly-4-hydroxybutyrate (P4HB), represents an alternative to standard allo- and xenograft implants, and are designed to combine synthetic materials' durability with biologics' biocompatibility
- There is a lack of comparative studies directly evaluating outcomes of biosynthetic vs. synthetic meshes in cancer patients

Objectives

Compare hernia recurrence and surgical risk factors in oncologic patients undergoing hernia repair with either biosynthetic or synthetic mesh reinforcement following prior abdominal or pelvic oncologic resection

Methods


Study population:
Patients > ≥ 18 years of age
Retrospective review, 2010-2023

Inclusion:
Prior history of abdominal or pelvic oncologic resection, which was complicated by hernia development and subsequently underwent repair with either biosynthetic or synthetic mesh

Results

Table 1. Operative Details			
Characteristic	Biosynthetic, N=103	Synthetic , N=109	p-value
ASA Class			0.5
II	6 (5.8%)	9 (8.3%)	
III	79 (77%)	86 (79%)	
IV	18 (17%)	14 (13%)	
Concomitant Urologic Procedure	8 (7.8%)	4 (3.7%)	0.2
Concomitant Gynecologic Procedure	14 (14%)	5 (4.6%)	0.022
Concomitant Large Bowel Procedure	16 (16%)	6 (5.5%)	0.017
Concomitant Small Bowel Procedure	7 (6.8%)	2 (1.8%)	0.094
Wound Class			0.011
Clean (I)	49 (48%)	75 (69%)	
Clean/Contaminated (II)	45 (44%)	29 (27%)	
Contaminated (III)	8 (7.8%)	4 (3.7%)	
Dirty/Infected (IV)	1 (1.0%)	1 (0.9%)	
Wound Class (Clean vs. Contaminated)			0.002
I	49 (48%)	75 (69%)	
II-IV	54 (52%)	34 (31%)	
Unilateral External Oblique Release	13 (13%)	15 (14%)	0.8
Bilateral External Oblique Release	71 (69%)	46 (42%)	<0.001
Unilateral Transverse Rectus Abdominus Release	0 (0%)	6 (5.5%)	0.029
Bilateral Transverse Rectus Abdominus Release	1 (1.0%)	8 (7.3%)	0.036
Depth of Mesh Placement			0.068
Onlay	22 (21%)	24 (22%)	
Retrorectus	70 (68%)	61 (56%)	
Underlay	11 (11%)	24 (22%)	
History of Ostomy	9 (8.7%)	5 (4.6%)	0.2
Ostomy at Time of Surgery	22 (21%)	9 (8.3%)	0.007

Table 2. Univariable Fine-Gray Competing Risk Regression Models for Reoccurrence with competing risk of death					
Characteristic	N	Event, N	HR ¹	95% CI ¹	p-value
Age at Surgery	212	16	1.01	0.96, 1.05	0.8
Body Mass Index	212	16	1.08	1.02, 1.14	0.005
Smoking Status					
Never Smoker	173	13	—	—	
Former Smoker	28	2	0.84	0.21, 3.37	0.8
Unknown Smoker Status	11	1	1.05	0.15, 7.55	>0.9
Radiation History	212	16	3.14	1.13, 8.71	0.028
Chemotherapy	212	16	0.77	0.29, 2.04	0.6
Prior Hernia Repair	212	16	1.53	0.54, 4.34	0.4
Mesh Degradation					
Biosynthetic	103	8	—	—	
Synthetic	109	8	1.17	0.45, 3.03	0.8
Wound Class,					
I	124	2	—	—	
II-IV	88	14	8.23	1.89, 35.8	0.005
Unilateral External Oblique	212	16	4.26	1.49, 12.2	0.007
Bilateral External Oblique	212	16	0.24	0.08, 0.75	0.014
Depth of Mesh Placement					
Onlay	46	9	—	—	
Retrorectus	131	2	0.09	0.02, 0.41	0.002
Underlay	35	5	0.81	0.27, 2.47	0.7
Ostomy at Time of Surgery	212	16	2.24	0.80, 6.27	0.12



Conclusion

- Biosynthetic & synthetic mesh are both effective options for ventral hernia repair in oncologic patients
- Retrorectus mesh placement and bilateral component separation are critical in reducing hernia recurrence risk in oncologic hernia repair
- Radiation increases the risk of hernia recurrence following repair

References

- Rosen MJ, Krpata DM, Petro CC, et al. Biologic vs synthetic mesh for single-stage repair of contaminated ventral hernias: a randomized clinical trial. JAMA surgery. 2022;157(4):293-301.
- Harris HW, Primus F, Young C, et al. Preventing recurrence in clean and contaminated hernias using biologic versus synthetic mesh in ventral hernia repair: the PRICE randomized clinical trial. LWW; 2021.
- Burger JWA, Luijendijk RW, Hop WCJ, Halm JA, Verdaasdonk EGG, Jeekel J. Long-term Follow-up of a Randomized Controlled Trial of Suture Versus Mesh Repair of Incisional Hernia. Annals of Surgery. 2004;240(4):578-585. doi:10.1097/01.sla.0000141193.08524.e7
- Song Y, Shannon AB, Roses RE, Fraker DL, Kelz RR, Karakousis GC. National trends in ventral hernia repairs for patients with intra-abdominal metastases. Surgery. Sep 2020;168(3):509-517. doi:10.1016/j.surg.2020.04.005