




The head and neck surgeon may be consulted for emergency surgical management of patients admitted for severe hypercalcemia due to primary hyperparathyroidism. Patient care must be integrated between hospitalists, the anesthesia team, and the managing surgeon. The crisis can be associated with secondary dyselectrolytemia and metabolic acidosis which can greatly complicate any planned surgical intervention. A thoughtful approach for fluid resuscitation can mitigate additional metabolic disturbances. The surgeon must be fully aware of the increased risk of metabolic perturbations, ectopic target locations, and parathyroid cancer in this patient subset.

Four patients from two senior authors that demonstrate a range of pathologies which can cause parathyroid crises are reported. The clinical presentation, laboratory data and preoperative imaging studies were reviewed. The medical resuscitation protocol, the surgical findings, final pathology and postsurgical course are summarized

Case	Ca ²⁺	PTH	Pathology
1	15.3	1258	Double adenoma
2	16.0	1371	Parathyromatosis vs. Parathyroid carcinoma
3	19.2	813	Retropharyngeal adenoma
4	12.2	587	Parathyroid carcinoma

Patient 1. Double Adenoma • Normal saline-induced hyperchloremic acidosis and hemodynamic instability		Patient 2. Parathyroiditis and mediastinal adenoma • 4 prior parathyroid surgeries	
Case Presentation 50 y.o. male admitted from ER w/ severe nausea & vomiting • Ca²⁺ 15.3 ; PTH 1258 Cr 3.46 • Hospitalization - Tx- volume repletion followed by normal saline / furosemide diuresis, pamidronate • Patient signed out AMA. Readmitted 8 days later: Recurrent nausea & vomiting • 9L urine output / 24 hours • nephrogenic diabetes insipidus due to hypercalcemia Urgent surgical preparation begins. Hypercalcemia treated w/ normal saline diuresis , calcitonin, and bisphosphonate with goal of Ca ²⁺ <12 prior to induction of anesthesia		Surgical Hx 1994: Bilateral exploration • R: 5x3 cm cystic mass adherent to thyroid extending from the superior pole into superior mediastinum 2010: Bilateral neck dissection • Ca ²⁺ > 19 ; PTH > 2000 • R: 2.1 cm cystic parathyroid mass • L: 1.5 cm cystic parathyroid mass • Biopsies and reimplanted into SCM 2011: Bilateral neck dissection • Removal of innumerable parathyroid nodules 2014: Tx with Sensipar • Ca ²⁺ > 12 2015: Bilateral neck dissection • > 20 tumors removed from R and L neck 2017: Bilateral exploration • Ca ²⁺ 16.0 ; PTH 903 • Mibi: uptake R neck and mediastinum • Two nodules firm with cicatrix → surgeon concerned for parathyroid cancer	
Surgery 		Operative findings Left inferior parathyroid adenoma • 4.2 x 2.4 x 1.2 cm • 9.9 grams Right superior parathyroid adenoma • 1.5 x 1.2 x 0.4 cm • 0.5 grams Pathology: hypercellular 1 year post-op • Ca ²⁺ 8.7 ; PTH 105 ; Cr 1.36	
2017 imaging Mibi scan: uptake in L lower neck and mediastinum 		Gamma probe identifying cervical parathyroid lesion 	
		Video assisted thoracic surgery showing mediastinal lesion 	
3 year post-op: Ca ²⁺ 9.1 ; PTH 326 ; Cr 2.0			

Case Presentation

69 y.o. male admitted w/ progressive lethargy

- Ca^{2+} 12.2 ; PTH 587
- Cr 2.2

Surgery

- Large tumor adherent to thyroid, extending into upper mediastinum
- Final path: parathyroid carcinoma



5 year post-op

- Ca^{2+} 8.8 ; PTH 39 ; Cr 1.95

Hypercalcemic crisis is a life-threatening condition most commonly due to primary hyperparathyroidism.¹ Prior to 1960, hypercalcemic crisis had a high fatality rate due to lack of prompt medical therapy.^{2,3}

Hyperparathyroidism crisis is characterized by a cycle of volume depletion leading to renal dysfunction/osmotic diuresis resulting in further volume contraction and hypercalcemia.⁴

The rate of calcium change determines the clinical presentation⁸:

- Renal dysfunction: polyuria, polydipsia (early), oliguric (late)
- GI symptoms: nausea, vomiting, constipation
- Neurologic symptoms: agitation, lethargy, coma

- Fluid resuscitation – Normal saline
 - Beware of hyperchloremic **metabolic acidosis**^{6,7}
 - **Consider alternative fluids**
- Lowering of serum calcium
 - Lasix, Bisphosphonate, Calcitonin
- Maximize renal function

Imaging considerations: 4D CT may be contraindicated due to renal failure

Surgical delay may result in worsening of hypercalcemic relapse and life-threatening complications. Consider surgical intervention at initial hospitalization.⁴

- Volume expansion and lowering calcium priority
- First report - normal saline resuscitation causing hyperchloremic acidosis subsequent to treatment of hypercalcemia
- Monitor chloride and bicarbonate
- Consider urgent surgery at same hospitalization
- Surgical considerations
 - Ectopic parathyroid adenomas
 - Multiglandular disease
 - Parathyromatosis
 - Parathyroid carcinoma

1. Ahmad S., Kuraganti G., Steenkamp. Hypercalcemic Crisis: A Clinical Review. Am. J. Med. 2015.
2. James P.R., Richards P.G. Parathyroid Crisis. Arch. Surgery. 1956;72(4):553-566.
3. Ziegler R. Hypercalcemic Crisis. J Am Soc Nephrol 83-89,2001.
4. Phitayakorn R., McHenry C.R. Hyperparathyroid Crisis: Use of Bisphosphonates as a Bridge to Parathyroidectomy. J Am Coll Surg 2008;206(3):1106-111
5. Kong H., Zhang, Z., Zhang, H. Anesthetic considerations in hyperparathyroid crisis - A case report. Medicine Jan 2021.100(1)
6. Scheingraber S. Rapid Saline Infusion produces hyperchloremic acidosis in patient undergoing gynecologic surgery. Anesthesia 1999 (90)
7. Gheorghe et al. Hyperchloremic Metabolic Acidosis Following Resuscitation of Shock. Chest 2010 Dec;238(6):2621-2.
8. Mousseaux C, et al. Epidemiology, clinical features, and management of severe hypercalcemia in critically ill patients. Ann Intensive Care. 2019 Nov 27;9(1):133.