

Morbidity and Mortality Following Total Hip and Knee Arthroplasty with Spinal vs. General Anesthesia: A Retrospective Analysis

Carlos Mucharraz, M.D¹., Samantha Diulus, B.S.², Daniel Schmitt, M.D.², Nicholas Brown, M.D.²

¹University of Colorado Anschutz Medical Campus, ²Loyola University Chicago Stritch School of Medicine



Background

Despite longstanding debate surrounding anesthesia in total joint arthroplasty, the topic remains controversial, and an optimal anesthetic mode has yet to be clearly identified. Historically, general anesthesia (GA) was recommended for total hip and knee arthroplasty (THA, TKA); however, in more recent years, spinal anesthesia (SA) has become more popular, especially in elderly patients with comorbidities [3]. Although multiple investigators have explored this topic, their conclusions are conflicting. Some studies have reported increased incidence of hypotension, motor blockade, urinary retention, and pruritis with SA [1,5,7,10]. On the other hand, some authors have reported increased blood loss, length of hospital stay, nausea, infection, and thromboembolic events with GA [4,8,9].

In comparison, hip fracture literature does not demonstrate superiority of SA or GA with multiple authors concluding that both methods are viable and comparable in morbidity and mortality [2,6,11]. This difference in reporting is interesting considering the similarities between THA and hip fracture repair. Therefore, despite previous studies examining anesthesia techniques in total joint arthroplasty, there is more room to explore this topic.

Objectives

The purpose of this study is to further investigate the relationship between total joint arthroplasty outcomes and anesthesia type in order to determine if there is a significant difference in morbidity and mortality between patients undergoing spinal anesthesia compared with those undergoing general anesthesia during primary total hip and knee arthroplasty.

Methods

Patients undergoing primary total hip or knee arthroplasty from February 2007 to February 2021 were retrospectively reviewed. From this population, two cohorts were created: GA (n = 2154) and SA (n = 3152). Readmission within 90 days, mortality within one year, and thromboembolic events in the postoperative 30 days were compared using logistic regression analysis. The multivariate model controlled for age and Charlson Comorbidity Index (CCI).

Results / Data

Table 1 demonstrates that there were several statistically significant differences in demographics between the two groups. Table 2 details that there was no difference in the odds of 90-day readmission or pulmonary embolism (PE) between the groups. The GA group demonstrated higher odds of experiencing a deep venous thrombosis (DVT) within 30 days postoperatively as well as higher odds of mortality within one year postoperatively in both the bivariate and multivariate logistic regression models.

Table 1. Demographics

	General Anesthesia n = 2154	Spinal Anesthesia n = 3152	p-value
Age (years)	63.51 ± 11.95 (20-94)	65.45 ± 10.35 (20-98)	< .001
BMI (kg/m2)	32.30 ± 8.71 (15.8-74.4)	32.50 ± 7.46 (15.2-64.1)	0.02
Sex, n (%)			0.441
Male	811 (37.7)	1221 (38.7)	
Female	1343 (62.3)	1931 (61.3)	
CCI	3.76 ± 2.82 (0-17)	3.87 ± 2.70 (0-18)	0.018

Values presented as mean ± SD unless otherwise indicated.
BMI: body mass index; CCI: Charlson Comorbidity Index; SD: standard deviation

Table 2. Postoperative Morbidity and Mortality According to Anesthesia Type

	General Anesthesia n = 2154	Spinal Anesthesia n = 3152	Bivariate logistic regression		Multivariate logistic regression	
			OR (95% CI)	p-value	OR (95% CI)	p-value
Thromboembolic Events						
DVT	69 (3.2)	58 (1.8)	1.77 (1.24-2.52)	0.002	1.66 (1.01 - 2.71)	0.044
PE	19 (.009)	18 (.006)	1.55 (0.81 - 2.98)	0.185	1.16 (0.51 - 2.59)	0.715
Readmission within 90 days	300 (13.9)	397 (12.6)	1.12 (0.96 - 1.32)	0.158	1.19 (0.98 - 1.44)	0.076
Mortality within 1 year	32 (.015)	20 (.006)	2.36 (1.36 - 4.21)	0.003	2.39 (1.00 - 5.79)	0.05

Values are presented as n (%).
OR: odds ratio; CI: confidence interval.
Odds ratio represents the odds of an event occurring in the GA group compared to the SA group.
Multivariate model was adjusted for age and CCI

Conclusion

Based upon this data, both spinal and general anesthesia are reasonable options for primary total joint arthroplasty with similar risk profiles. However, general anesthesia may be associated with a higher rate of DVT.

References

- 1.Davis, F. M. Laurenson, V. G., Gillespie, W. J., Wells, J. E., Foote, J., & Newman, E. (1989). Deep vein thrombosis after total hip replacement. A comparison between spinal and general anaesthesia. *The Journal of bone and joint surgery: British volume*, 71(2), 181–185.
- 2.Guay, J., Parker, M. J., Gajendragadkar, P. R., & Kopp, S. (2016). Anaesthesia for hip fracture surgery in adults. The Cochrane database of systematic reviews, 2(2), CD000521.
- 3.Indelli, P. F., Grant, S. A., Nielsen, K., & Vail, T. P. (2005). Regional anesthesia in hip surgery. *Clinical orthopaedics and related research*, 441, 250–255.
- 4.Maurer, S. G., Chen, A. L., Hiebert, R., Pereira, G. C., & Di Cesare, P. E. (2007). Comparison of outcomes of using spinal versus general anesthesia in total hip arthroplasty. *American journal of orthopedics (Belle Mead, N.J.)*, 36(7), E101–E106.
- 5.Modig, J., Borg, T., Karlström, G., Maripuu, E., & Sahlstedt, B. (1983). Thromboembolism after total hip replacement: role of epidural and general anesthesia. *Anesthesia and analgesia*, 62(2), 174–180.
- 6.Morgan, L., McKeever, T. M., Nightingale, J., Deakin, D. E., & Moppett, I. K. (2020). Spinal or general anaesthesia for surgical repair of hip fracture and subsequent risk of mortality and morbidity: a database analysis using propensity score-matching. *Anaesthesia*, 75(9), 1173–1179.
- 7.Pugely, A. J., Martin, C. T., Gao, Y., Mendoza-Lattes, S., & Callaghan, J. J. (2013). Differences in short-term complications between spinal and general anesthesia for primary total knee arthroplasty. *The Journal of bone and joint surgery: American volume*, 95(3), 193–199.
- 8.Pu, X., & Sun, J. M. (2019). General anesthesia vs spinal anesthesia for patients undergoing total-hip arthroplasty: A meta-analysis. *Medicine*, 98(16), e14925.
- 9.Park, Y. B., Chae, W. S., Park, S. H., Yu, J. S., Lee, S. G., & Yim, S. J. (2017). Comparison of Short-Term Complications of General and Spinal Anesthesia for Primary Unilateral Total Knee Arthroplasty. *Knee surgery & related research*, 29(2), 96–103.
- 10.Rashiq, S., & Finegan, B. A. (2006). The effect of spinal anesthesia on blood transfusion rate in total joint arthroplasty. *Canadian journal of surgery: Journal canadien de chirurgie*, 49(6), 391–396.
- 11.Shelton, C., & White, S. (2020). Anaesthesia for hip fracture repair. *BJA education*, 20(5), 142–149.