

Treatment of Pain Using Cervical Radiofrequency Ablation: Outcomes from an International, Prospective Multicenter Study (RAPID)

Joseph Atallah¹, Bradley Holt², Michael Danko³, David Provenzano⁴, Maaz Iqbal⁵, Binit Shah⁶, Albert Singh⁷, Harsh Sachdeva⁸, Ella Ver Donck⁹, Bart Liebrand¹⁰, Erik Shaw¹¹, Sherri Haas¹², Rajat Sekhar¹³, Ann Pan¹⁴, Edward Goldberg¹⁴

1. The Toledo Clinic, Toledo, OH USA 2. Tucson Orthopaedic Institute, Tucson, AZ USA 3. Premier Pain Treatment Institute, Loveland OH, USA 4. Pain Diagnostics and Interventional Care, Sewickley, PA USA 5. Elite Pain and Spine Institute, Mesa, AZ USA

6. Carolinas Pain Center, Huntersville, NC USA7. Quincy Medical Group, Quincy, IL USA 8. UC Health Pain Medicine, Cincinnati, OH USA 9. AZ Sint-Jan, Bruges, Belgium 10. St Anna Hospital, Geldrop, The Netherlands

11. Shepherd Center, Shepherd Pain Institute, Atlanta, GA, USA 12. Twin Cities Pain Clinic & Surgery Center, Edina, MN USA 13. St. Vincent’s Medical Center, Bridgeport, CT USA 14. Boston Scientific Neuromodulation, Valencia, CA USA

BACKGROUND

Chronic axial neck pain with or without headache and/or upper extremity pain is increasingly noted as a source of enduring disability and medical costs, and one of the leading causes of healthcare burden.¹ Several clinical studies, including RCTs and observational studies have demonstrated positive outcomes using radiofrequency ablative (RFA) techniques for a range of chronic neck pain indications.²⁻⁶ Further, a recent meta-analysis of 20 previously published studies showed that evidence collectively published supports the use of therapeutic radiofrequency (RF) techniques for the effective management of chronic neck pain.⁷ In light of the on-going opioid epidemic and the need to provide better care to chronic pain patients, recent guidelines (HHS 2019 Best Practices report and 2022 CDC guidelines) have highlighted the need to incorporate interventional techniques including neuromodulation and RF.⁸⁻¹⁰ Thus, it is important to continually assess and document clinical effectiveness of contemporary RF devices in real-world patients with chronic neck pain, headaches and proximal upper extremity pain. We assessed patients diagnosed with chronic cervical facet joint pain treated with RFA in a prospectively-enrolled, real-world, multicenter, international study.

METHODS

Study Design	Multicenter, Prospective, International Outcomes Study RAPID Study (clinicaltrials.gov identifier: NCT04673032)
Study Device	Commercially-approved RFA Systems (Boston Scientific)
Subjects	•75-enrolled patients with cervical facet joint pain •73-treated patients with cervical facet joint pain (initial procedure completed)
Study Eligibility Criteria	Key Inclusion Criteria: Study candidate is scheduled to be treated with a commercially approved Boston Scientific RF system for pain per local Directions for Use (DFU) Key Exclusion Criteria: Meets any contraindications per locally applicable Directions for Use (DFU)

STUDY SCHEMATIC

Informed Consent

Baseline Visit

Day 0: Procedure Visit

1 mo. post-procedure

3 mos. post-procedure

6 mos. post-procedure

12 mos. post-procedure

18 mos. post-procedure

24 mos. post-procedure

Study Completion

RESULTS

Baseline Characteristics
(n = 73 patients with initial RFA completed)

Gender - Females (%)	60.3% (n = 44/73)
Age (years) [Mean (SD)]	59.6 ± 11.3 years (n = 73)
Pain Duration (years) [Mean (SD)]	13.7 ± 11.5 years (n = 73)
Baseline Targeted Pain VRS Score [Mean (SD)]	VRS: 6.62 ± 1.6
Number of Study RF Procedures [Mean (SD)]	2.3 ± 1.4 procedures (n = 73)
Follow-up Duration [Mean (SD)]	498.5 ± 235.4 days (n = 73)
Location Targeted by RFA	Head, neck, shoulder and upper back – 75.3% (55/73) Mid/Low Back – 24.7% (18/73)
Pain Type Treated by RFA	Cervical zygapophyseal joints pain – 100% (73/73)

Targeted Pain is the area of pain intended to be treated with RF

Targeted Pain Scores up to 24-months post-procedure

VRS Scores (0-10)

Baseline (n = 73) 3-month follow-up (n = 68) 6-month follow-up (n = 61) 12-month follow-up (n = 48) 24-month follow-up (n = 20)

6.62 2.62 2.93 3.48 2.8

Δ = 3.4 (p<0.0001)

•At 3-months, a mean 3.9-point reduction in VRS score was observed (p<0.0001)

•At 24-months, a mean 3.4-point reduction in VRS score was observed (p<0.0001)

PPR = percent pain relief

Patient Satisfaction (PGIC) post-procedure

Patient Satisfaction - Percent Improvement

3-month follow-up (n = 67) 6-month follow-up (n = 61) 12-month follow-up (n = 48) 24-month follow-up (n = 20)

25.4% 23.0% 27.1% 10.0%

32.8% 44.3% 35.4% 40.0%

35.8% 24.6% 25.0% 40.0%

4.5% 6.5% 4.2% 10.0%

no change very much improved much improved minimally improved

~90% of patients reported improvement (very much, much, or minimally improved) up to 24-months post-procedure

Responder Rates post-procedure

Proportion Reporting %

3-month follow-up (n = 68) 6-month follow-up (n = 61) 12-month follow-up (n = 48) 24-month follow-up (n = 20)

79.4% 72.1% 71% 85%

89.7% 90.2% 85.4% 85%

50% or more improvement 30% or more improvement

At all timepoints:
•≥85% responder rate (≥30% overall PPR)
•≥71% responder rate (≥50% overall PPR)

CONCLUSIONS

•Preliminary sub-analysis derived from this ongoing, prospective, multicenter, real-world clinical outcomes RF study of 73 patients with chronic cervical pain demonstrates significant improvement in pain scores out to 24-months post-procedure.

•Follow-up outcomes:
□3.4-point VRS pain score improvement out to to 24-months post-procedure
□Responder rate ≥71% (≥50% overall PPR) and ≥85% (≥30% overall PPR) at all time points
□~90% of patients reported improvement (i.e., very much, much or minimally improved)

REFERENCES

1.Murray CJ, Atkinson C, Bhalla K, et al. The state of US health, 1990-2010: burden of diseases, injuries, and risk factors. JAMA. 2013 Aug 14;310(6):591-608.

2.Lord SM, Barnsley L, Wallis BJ, Bogduk N. Chronic cervical zygapophyseal joint pain after whiplash. A placebo-controlled prevalence study. Spine (Phila Pa 1976). 1996 Aug 1;21(15):1737-44; discussion 1744-5.

3.Barnsley L. Percutaneous radiofrequency neurotomy for chronic neck pain: outcomes in a series of consecutive patients. Pain Med. 2005 Jul-Aug;6(4):282-6.

4.Manchikanti L, Singh V, Falco FJ, Cash KA, Fellows B. Comparative outcomes of a 2-year follow-up of cervical medial branch blocks in management of chronic neck pain: a randomized, double-blind controlled trial. Pain Physician. 2010 Sep-Oct;13(5):437-50.

5.MacVicar J, Borowczyk JM, MacVicar AM, Loughnan BM, Bogduk N. Cervical medial branch radiofrequency neurotomy in New Zealand. Pain Med. 2012 May;13(5):647-54.

6.Speldewinde GC. Outcomes of percutaneous zygapophyseal and sacroiliac joint neurotomy in a community setting. Pain Med. 2011 Feb;12(2):209-18.

7.Manchikanti L, Knezevic NN, Knezevic E, et al. A Systematic Review and Meta-analysis of the Effectiveness of Radiofrequency Neurotomy in Managing Chronic Neck Pain. Pain Ther. 2022 Nov 24:1-48.

8.Dowell D, Ragan KR, Jones CM, Baldwin GT, Chou R. CDC Clinical Practice Guideline for Prescribing Opioids for Pain - United States, 2022. MMWR Recomm Rep. 2022 Nov 4;71(3):1-95

9.U.S. Department of Health and Human Services (2019, May). Pain Management Best Practices Inter-Agency Task Force Report: Updates, Gaps, Inconsistencies, and Recommendations. Retrieved from U. S. Department of Health and Human Services website: <https://www.hhs.gov/ash/advisory-committees/pain/reports/index.html>

10.Hurley RW, Adams MCB, Barad M, Bhaskar A, Bhatia A, Chadwick A, Deer TR, Hah J, Hooten WM, Kissoon NR, Lee DW, McCormick Z, Moon JY, Narouze S, Provenzano DA, Schneider BJ, van Eerd M, Van Zundert J, Wallace MS, Wilson SM, Zhao Z, Cohen SP. Consensus practice guidelines on interventions for cervical spine (facet) joint pain from a multispecialty international working group. Reg Anesth Pain Med. 2022 Jan;47(1):3-59.

DISCLOSURES

Study Sponsored by Boston Scientific.
Drs. Atallah and Provenzano have consulting agreements with Boston Scientific. Ann Pan and Edward Goldberg are employees of Boston Scientific

ASRA
PAIN MEDICINE

23rd Annual Pain Medicine Meeting

November 21-23, 2024
Caesars Palace | Las Vegas, Nevada
#ASRAFall24