

Hydrogen fuel production on MoS₂ nanotubes



Charlie Ruffman, J. T. A. Gilmour, Anna L. Garden*

1 Motivation

H₂ is a promising carbon-zero fuel, which can be produced cleanly by the **electrolysis** of water, using a **catalyst**.^[1]


MoS₂ nanosheets are a promising hydrogen evolution catalyst. However, the flat basal plane – which makes up most of the material – is inactive.

In this density functional theory study, we examine how coiling MoS₂ into **nanotubes** can enhance catalysis.




2 What makes a good catalyst?

H⁺ must (1) adsorb to the catalyst, (2) combine with another H, then (3) desorb as H₂.^[2] Density functional theory (BEEF-vdW, VASP) is used to calculate the energetics:



If a catalyst adsorbs H too strongly ($\Delta G_{H,ads} < 0$) or weakly ($\Delta G_{H,ads} > 0$), an **energy cost** must be paid to either get H onto the surface or remove it again after reaction.

A **ideal catalyst** both adsorbs and desorbs H without an energy penalty ($\Delta G_{H,ads} = 0$).^[3] The closer $\Delta G_{H,ads}$ is to zero, the better the catalyst.




Want to
know
more?



Ruffman, C.; Gilmour, J. T. A.;
Garden, A. L. *Nanoscale
Advances* 2021.
DOI: 10.1039/d1na00441g

3 MoS₂ nanotubes


Pristine nanotubes:



$\Delta G_{H,ads}$ on nanotubes is **lowered** relative to flat MoS₂, especially at small diameters.



At S-vacancy defect sites, $\Delta G_{H,ads}$ is **close to thermoneutral** (0 eV) when the nanotubes are 18–20 Å in diameter. These materials may make very good hydrogen evolution catalysts!

S-vacancy defects:




What causes size-dependent tuning of H adsorption?

As the nanotubes get smaller, the Mo d-states get closer in energy to the H s-states. This leads to a **stronger bond** between MoS₂ and H.^[4]



For the S-vacancy defect site, we find a reliable linear relationship between $\Delta G_{H,ads}$ and the Mo d-state to H s-state energy gap.

Controlling the size of MoS₂ nanotubes offers a clear way to **fine-tune and improve** their hydrogen evolution activity.



References

- [1] Jaramillo, T.; Jørgensen, K.; Bonde, J.; Nielsen, J.; Hørch, S.; Chorkendorff, I. *Science* 2007, 317, 100-102.
- [2] Ruffman, C.; Gordon, C. K.; Skúlason E.; Garden, A. L. *J. Phys. Chem. C* 2020, 124, 17015-17026.
- [3] Tsai, C.; Chan, K.; Nørskov, J. K.; Abild-Pedersen, F. *Surf. Sci.* 2015, 640, 133-140.
- [4] B. Hammer B.; Nørskov, J. *Surf. Sci.* 1995, 343, 211–220.

@CharlieRuffman

*MacDiarmid Institute for Advanced Materials
and Nanotechnology & University of Otago.
New Zealand