

Effective Photocatalytic Seawater Splitting



Rohit Mann and Deepa Khushalani
Department of Chemical Science, Tata Institute of Fundamental Research, Mumbai

Abstract


The photocatalytic (PC) hydrogen production from water splitting is a promising route to fulfill the current energy demand in an eco-friendly and sustainable manner. For photocatalysis to become industrially viable, seawater should be used. This solvent owes its salinity due to various ions (Na^+ , K^+ , Mg^{2+} , Ca^{2+} , Cl^- , Br^- , SO_4^{2-} , and CO_3^{2-}). To date, there are contradicting reports in the literature on the influence of these ions on the PC splitting of water. A variety of different neutral transition metal-oxide-based photocatalysts has been exploited. Although they might be effective in splitting deionized water^[1], however, thus far, there has been a lack of a well-established route to effectively using any semi-conductor for seawater splitting. Presented here are the details of our work where an intriguing, ionized, low-cost semiconductor has been successfully evaluated for sea-water splitting. The structure consists of a modified carbon nitride such that it forms an ionized organic polymer-based system.^[2] A detailed study has been done using this salt-type semiconductor in the presence of various ions (Na^+ , K^+ , Mg^{2+} , Ca^{2+} , Cl^- , SO_4^{2-}), and their role has been probed in modulating the photocatalytic activity. Photoelectrochemical measurements have provided insight as to how the presence of cations aids advantageously in forming an effective *in-situ* interface between catalyst/co-catalyst for improved charge transfer. This improved interfacial charge transfer rationalizes the 8-fold enhancement in the photocatalytic rate in the presence of simulated seawater compared to deionized water. It provides an impetus for using these carbon nitride structures for sustainable PC splitting of seawater.

Introduction


	Fuel	Specific Energy density (MJ/Kg)
Fossil Fuel	Hydrogen gas	141
	Natural gas	53.6
	Gasoline	46.4
	Diesel	45.6
Capacitor	Supercapacitor	0.018
Battery	Lithium ion battery	0.87
	Nickel metal hydride battery	0.41

Current Usage of Hydrogen


Annually approx. > 800 metric tons of CO_2 emitted into the environment to generate hydrogen only



Synthesis



Photocatalytic Water Splitting




Only the cationic part of the electrolyte plays a role in augmenting the H_2 production.

➤ Hydrogen is having the highest specific energy density


➤ Combustion of hydrogen has zero carbon emission

Photo-catalysis



Photocatalyst	Cocatalyst	H_2 Production $\mu\text{mole h}^{-1} \text{g}^{-1}$	Seawater type	Better than Pure water?	% Change in Performance
TiO_2 [3]	CuO	3.1	3% NaCl	No	↓33%
p-GaN/InGaN [4]	Rh/Cr ₂ O ₃ , cobalt oxide	5.1×10^6	Simulated seawater	Yes	↑35%
$\text{La}_2\text{Ti}_2\text{O}_7$ [5]	Ni	69.6	Natural seawater	No	↓49%
$\text{WO}_2\text{-Na}_2\text{WO}_3$ [6]	-	5.76	Seawater	Yes	↑15%
$\text{ZnS}_{1-x}\text{-xO}_{0.5}\text{O}_2(\text{OH})_x\text{-ZnO}$ [7]	NiS	354	Simulated water	Yes	↑11%
Covalent organic polymer COP-TP [8]	Pt	4250	Simulated water	No	↓15%
CdS nanosheets [9]	Carbon dots (CDs)	4640	Natural Seawater	No	↓54%

Do these ions affect the rate of photocatalytic water splitting?



☐ Metal-free photocatalyst


☐ Visible absorption cross-section

☐ Proper band alignment for water splitting

☐ Stable in acidic and basic conditions


☐ Ease of preparation, inexpensive

Characterization




Photodeposition of Pt cocatalyst removes the negative charged from the framework

The formation of blue species in DI water and K_2SO_4 show that electrons are stored on the material



Reduction in photocurrent with Bi-valent cation showcases an effective interface formed between catalyst/cocatalyst.

Conclusion & Future Direction



☐ The augmentation in photocatalysis for ionized structure is substantially high compared to neutral transition metal oxide.

☐ The Presence of cations improves the electron transfer ability from the catalyst to the cocatalyst.

☐ Ionized carbon nitride showcases the promising behavior for seawater splitting.

Reference

- [1] Jining Zhang *et al.* *Nano Research*, 2020, 13(9), 2313–2322
- [2] Tanmay Banerjee *et al.* *Nat. Rev. Mater.*, 2020, 6, 168–190.
- [3] Simamora A. J. *et al.* *Int. J. Hydrog. Energy* 2012, 37, 18, 13855–13858
- [4] Guan X. *et al.* *J. Phys. Chem. C* 2018, 122, 13797–13802
- [5] Ji S. M. *et al.* *J. Photochem. Photobiol. A* 2007, 189, 141–144
- [6] Cui G. *et al.* *Nano. Lett.* 2015, 15, 7199–7203
- [7] Li Y. *et al.* *Int. J. Hydrog. Energy* 2011, 36, 10565–10573
- [8] Liu Y. *et al.* *ACS Appl. Mater. Interfaces* 2018, 10, 30698–30705
- [9] Zhu C. *et al.* *Appl. Catal. B.* 2019, 242, 178–185

Acknowledgment

Prof. Vivek Polshettiwar and his lab members Dr. Sushma Rawal and Rajesh Dr. Malay Patra, Devidas Ji, and Dr. Subhankar for ICP-MS measurements My lab members, Balasaheb B Chandanshive, Ashish, Payal, Komal, Subhra And DAE for Funding

How an ionized carbon nitride photocatalytic water splitting is going to modulate in the presence of ions?

XPS confirms F-PCN has a negatively charged framework