


Synthesis, immobilization and characterization of palladium complexes for Suzuki-Miyaura coupling reactions

Mosiua, M.E., Dr. Mahamo, T. & Prof. Sekota, M.



NATIONAL UNIVERSITY OF LESOTHO



Affiliations:


INTRODUCTION

The Palladium catalyzed Suzuki-Miyaura cross-coupling reaction is one of the most reliable methods for the construction of carbon-carbon bonds in solution. Herein, the synthesis and characterization of palladium complexes containing salicylaldehyde and iminophosphine ligands are reported. The ligands are prepared conveniently in high yield via Schiff base condensation reactions between salicylaldehyde with appropriate primary amines for salicylaldehydes and 2-diphenylphosphinobenzaldehyde and the same amines used for the iminophosphine ligands. The ligands were subsequently used to synthesize the palladium complexes of the type $[Pd(P^N)Cl_2]$ and $[Pd(O^N)Cl_2]$ from metal precursor $[Pd(cod)Cl_2]$. The complexes were isolated as air stable solids. The complexes were immobilized onto magnetite (Fe_3O_4) coated chicken feathers giving $[Pd(P^N)Cl_2]-CF@Fe_3O_4$ and $[Pd(O^N)Cl_2]-CF@Fe_3O_4$ complexes, which will behave as heterogeneous catalysts.

OBJECTIVES


- To synthesize and characterize iminophosphine and salicylaldehyde Palladium complexes
- To immobilize the synthesized complexes
- To test the catalysts' activity in Suzuki-Miyaura coupling reactions
- To determine recyclability and reusability of the catalysts

EXPERIMENTAL APPROACH




Scheme 1: synthesis of salicylaldehyde ligands


EXPERIMENTAL APPROACH (Cont...)




Scheme 2: Synthesis of iminophosphine ligands



Scheme 3: complexation of synthesized ligands using $Pd(COD)Cl_2$




Scheme 4: Synthesis of magnetite nanoparticles



Scheme 5: immobilization process of synthesized complexes onto chicken feather coated magnetite

Typical Suzuki-Miyaura Cross-Coupling Reaction



Scheme 7: catalysis – Suzuki-Miyaura coupling reaction

RESULTS

Table 1: List of primary amines used for both salicylaldehyde and iminophosphine ligands

Primary amine
Ethylenediamine
2-Thiophenemethylamine
2-Picolylamine
4-Aminobenzoic acid
Furfurylamine




Figure 1: some of the ligands synthesized




Figure 2: ^1H -NMR spectrum of salen ligand

CONCLUSION & FUTURE WORK

5 salicylaldehyde ligands and their complexes were synthesized and immobilized onto magnetite coated chicken feathers

2 iminophosphine ligands and their complexes were synthesized and coated onto magnetite coated chicken feathers

Still awaiting characterization results – NMR, IR, XRD, BET of the samples from N.W.U

Catalytic tests run but awaiting GC results from N.W.U

Further work needed on iminophosphine ligands and complexes, catalytic studies to have a constructive comparable results and test the recyclability and reusability of the catalysts.

Analysis and outline the impact of magnetite and chicken feathers on the complexes, and ligands using Secondary Coordination Sphere analogy.

REFERENCES

- T. Mahamo *et al.*, "Neutral palladium (II) complexes with P , N Schiff-base ligands : Synthesis , characterization and application as Suzuki e Miyaura coupling catalysts," *J. Organomet. Chem.*, vol. 703, pp. 34–42, 2012, doi: 10.1016/j.jorganchem.2011.12.021.
- S. Y. Abbas, "Synthesis , characterization and antimicrobial activity of 5 - (arylazo) salicylaldehydes and their copper (II) complexes," no. July 2017, pp. 1–10, 2018, doi: 10.1002/aoc.4032.
- R. Rahmawati, A. Taufiq, S. Sunaryono, A. Fuad, and B. Yulianto, "Synthesis of Magnetite (Fe_3O_4) Nanoparticles from Iron sands by Co- precipitation- Ultrasonic Irradiation Methods," vol. 9, no. 1, pp. 155–160, 2018.

ACKNOWLEDGEMENTS & CONTACTS

I would like to thank both my supervisors, Dr. Mahamo & Prof. Sekota for the guidance, and the University of Lesotho & North-West University Chemistry departments support.

edlettmosiua@gmail.com