




Nanoscale Celecoxib Prodrugs: As efficient Anti-Inflammatory Principles



Rudaba Saleem^{ab}, Mujeeb-ur-Rehman^{ab}, Saadat Hussain^a, Almas Jabeen^c

^aH.E.J. Research Institute of Chemistry, International Center for Chemical and Biological Sciences, University of Karachi, Karachi-75270, Pakistan; Universite Paris-Saclay, institlute 91400 Orsay, France. Dr. Pnajwani Center for Molecular Medicine and Drug Research, International Center for Chemical and Biological Sciences, University of Karachi, Karachi-75270

Rheumatoid Arthritis




Abstract

Celecoxib Prodrugs were synthesized with three different biodegradable linkers, and their structures were confirmed by various spectroscopic techniques. Prodrugs 1-3 were encapsulated into Nanoparticles via the sole addition of DSPE-PEG₂₀₀₀. Physicochemical characterization of these NPs was analyzed. Further, these NPs were subjected to in vitro studies and found active against cytokines like TNF- α , MCP-1 and IL- β . In Vivo studies reveal NPs are potent but comparatively less active than celecoxib


Introduction

- Celecoxib is a non-steroidal anti-inflammatory drug (NSAID) that selectively inhibits cyclooxygenase-2 (CoX-2) to alleviate inflammation.
- Despite its therapeutic benefits, it faces challenges, including low bioavailability, unfavorable pharmacokinetics, and potential cardiac toxicity.
- To enhance its efficacy, we have developed nanoparticles designed to improve hydrophobicity, biodegradability and stability profile.


Graphical Abstract




Synthesis and Characterization of Prodrugs (1-3)




Synthesis of Nanoparticles (Emulsion Evaporation)




Physicochemical Characterization of Nanoparticles



In-Vitro Studies



In-Vivo Studies (Prodrug-3 NPs)



Conclusion

We have successfully synthesized prodrugs of celecoxib with different biodegradable linkages using the C16 chain (1-3). Furthermore, we have synthesized nanoparticles with DSPE-PEG2000, which have an average size of 150 nm and a zeta potential of -32.5 mV. The encapsulation efficiency has been found to be 99%. In in vitro studies, these nanoparticles have shown activity against inflammatory cytokines. In in vivo studies, Prodrug-3 Nps have exhibited some activity against CIA mice, but they are not as effective as the free drug.

Future Prospects

- We are set to validate prodrugs 1 and 2 against CIA model mice.
- We plan to effectively design prodrugs that ensure sustained and controlled release in biological media with a detailed pharmacokinetics profile.

References

Saleem, R., Rehman, M. U., Hussain, S., Arif, A., Malik, H. N., Naqvi, F., & Jabeen, A. (2023). Nanoscale celecoxib prodrugs: As efficient anti-inflammatory principles. *Journal of Drug Delivery Science and Technology*, 90, 105089.

Acknowledgment

We would also like to acknowledge the organizers of Pakistan-France PERIDOT Research Mobility Program. No.PhaseVII-1/PERIDOT.R&D/HEC/2021), jointly implemented by the Higher Education Commission, Pakistan, and Ministry of Foreign Affairs & International. Development (MAEDI) and the Ministry of Higher Education and Research (MESRI), France.