

POST-SURGICAL DIABETIC FOOT ULCERS:

Outcomes with Injectable Sodium Hyaluronate and Collagen Amino Acids in Clinical Practice

Marco Mastinu, MD¹; Federico Rolando Bonanni, MD²; Luigi Uccioli, MD, Professor²; Davide Lauro, MD, Professor²; Marco Meloni, MD, Professor²
¹S.C. di Diabetologia e Malattie Metaboliche, ASL5 – Oristano; ² University of Tor Vergata, Roma.

Dual-center retrospective single-arm study • 12-week follow-up • 10 patients

BACKGROUND & OBJECTIVE

Deep post-surgical Diabetic Foot Ulcers (DFUs) are frequently associated with delayed healing and a high risk of complications. This study evaluated the efficacy and safety of weekly injectable sodium hyaluronate combined with collagen amino acids, added to standard of care


METHODS

- Dual-center, retrospective, single-arm study
- Patients with post-surgical DFUs
- Weekly injectable treatment + standard of care
- Secondary neutral dressing
- Follow-up: 12 weeks

OUTCOMES

Complete healing • Time to healing • ≥50% wound reduction • Granulation • Infection • Adverse events • Surgical re-intervention


PATIENT PROFILE



BASELINE ULCER SIZE

7.2 ± 2.3 cm²


KEY CLINICAL OUTCOMES



CONCLUSIONS

Injectable Sodium Hyaluronate combined with Collagen amino acids showed a favorable safety profile and promising clinical outcomes. Controlled studies are needed to confirm these preliminary findings

CASE - 1



CASE - 2

