

Comprehensive clinical and laboratory assessment of visceral leishmaniasis (VL) patients

ELENI AYELE, SASKIA VAN HENTEN, ASNAKEW ENGIDAW, TIGIST MEKONNEN, TADFE BOGALE, ANNELIES MONDELAERS, SAÏD ABDELLATI, MEKIBIB KASSA, MYRTHE PAREYN, JOHAN VAN GRIENSVEN, CPS STUDY TEAM

Leishmaniasis Research and Treatment Center, University of Gondar Hospital, Gondar, Ethiopia; Institute of Tropical Medicine Antwerp, Belgium

BACKGROUND

- **Ethiopia** has the **second highest burden** of VL disease among Sub-Saharan African countries.
- Many studies, including clinical trials, focus on a selected patient population or marker
- **Poor treatment outcomes** for VL have become a challenging public health problem.
- **Relapse** is increasingly recognized as an important clinical challenge, with 10-15% estimated to develop relapse in the Ethiopian setting.

OBJECTIVES

GENERAL OBJECTIVE

To **systematically** describe clinical symptoms, investigations, laboratory tests and outcomes amongst VL patients

SPECIFIC OBJECTIVES

- To analyze treatment outcomes (**relapse, treatment failure** and **death**)
- To determine **how many VL patients have tuberculosis** (based on chest Xray, GeneXpert, ultrasound, and TB-LAM)
- To describe **blood culture results**
- To describe **antibiotic prescription patterns** and whether the CRP and other investigations support antibiotic prescription
- To describe **electrolyte and INR disturbances** before and during treatment

METHODS

- **Prospective** cohort study in *Leishmania* Research and Treatment Center (**LRTC**), University of Gondar Hospital. An average of 300 to 400 patients treated for VL annually.
- Study participants (≥ 12 years) with **clinically or parasitologically confirmed VL** will be recruited.
- Total estimated study recruitment period will be around two years.
- Patients are followed closely **during treatment and follow-up visits until 12 months after end of treatment for checking late relapse.**
- Status: recruitment started in FEB 2023. Currently, around **200 patients are enrolled**, and data entry is ongoing.




FIG 1. STUDY SCHEDULE

PRELIMINARY RESULTS FOR 52 PATIENTS

- 44/52 patients had primary VL, 8 patients had relapse VL
- 12/52 patients were coinfectd with HIV
- 6/52 patients had a TB coinfection, of which 2 were diagnosed during VL treatment
- Ingesting/inhaling traditional medication was common (21/52). Many such patients had hepatotoxicity with elevated transaminases, INR, and signs of liver failure (**Fig 2**)
- SSG/PM was most commonly given, 8 patients were started on AmBisome, 12 HIV patients received AmBisome/Miltefosine

FIG 2. VL Patient who died due to liver failure having ingested traditional treatment





FIG 3. EoT Treatment outcomes



- Antibiotics were frequently given (39/52), and were mostly from the Watch category
- Median CRP results at baseline were 32 (range 2-190)
- 25/52 patients had hypokalemia at D0
- 3 patients developed PKDL during follow-up




FIG 4. PCR Results shown in Ct-values by end of treatment outcome and HIV status.