

# Quality of routine data related to facility-based maternal mortality measurement in Kampala City, Uganda

Birabwa C<sup>1,2,3\*</sup>, Banke-Thomas A<sup>4</sup>, Semaan A<sup>2</sup>, van Olmen J<sup>3</sup>, Kananura RM<sup>1</sup>, Arinaitwe ES<sup>5</sup>, Waiswa P<sup>1,6,7</sup>, Beňová L<sup>2</sup>

<sup>1</sup>Department of Health Policy, Planning and Management, Makerere University School of Public Health, Kampala, Uganda <sup>2</sup>Department of Public Health, Institute of Tropical Medicine, Antwerp, Belgium <sup>3</sup>Department of Family Medicine and Population Health, University of Antwerp, Antwerp, Belgium <sup>4</sup>Faculty of Epidemiology and Population Health, London School of Hygiene and Tropical Medicine, London, United Kingdom <sup>5</sup>Minsistry of Health, Kampala Uganda <sup>6</sup>Global Public Health, Karolinska Institute, Stockholm, Sweden <sup>7</sup>Busoga Health Forum, Jinja, Uganda  
\*[cathybirabwa3@gmail.com](mailto:cathybirabwa3@gmail.com)

DOI: [10.21203/rs.3.rs-3528370/v1](https://doi.org/10.21203/rs.3.rs-3528370/v1)

## BACKGROUND AND OBJECTIVE

Regular quality assessments are necessary to improve the reliability of routine data for different purposes, including estimating facility-based maternal mortality<sup>1,2</sup>. This study sought to assess the quality of routine data on deliveries, livebirths and maternal deaths in Kampala City, Uganda.


## METHODS

**Design:** Desk review of routine data using the World Health Organization data quality review framework

**Data:** From all facilities that reported at least one delivery between 2016 and 2019, and a subset of four facilities designated to provide comprehensive obstetric care


**Outcome:** Data completeness and consistency

**Analysis:** Descriptive, generated performance for whole dataset and for facilities at health centre (HC) III and higher levels



## RESULTS

- Prevalence of outliers was generally <2% for the three data elements
- Inconsistencies over time were mostly observed for maternal deaths
- Data from HC III and higher-level facilities had better quality metric scores compared to lower-level facilities.



**References:** <sup>1</sup>Mallick L, Gheda T, Sorrel N, Trinadh D, Wenjuan W. Using Health Management Information Systems Data to Contextualize Survey-Based Estimates of Fertility, Mortality, and Wasting. DHS Occasional Paper No. 12. Rockville, Maryland, USA; 202. <sup>2</sup>Agiraembabazi G, Ogwal J, Tashobya C, Kananura RM, Boerma T, Waiswa P. Can routine health facility data be used to monitor subnational coverage of maternal, newborn and child health services in Uganda? BMC Health Serv Res. 2021 Sep 1;21

## CONCLUSION

Routine data from facilities that provided childbirth services in Kampala (2016-2021) were of better quality in facilities that routinely conduct deliveries. Given likely underreporting of maternal deaths, further efforts to verify and count all maternal deaths in facilities are essential for more reliable estimates.