



POSTPARTUM DEPRESSION

AMONG WOMEN AFTER GIVING BIRTH IN A MOUNTAINOUS AREA OF VIETNAM

Thi Thanh Huong NGUYEN¹; Manh Linh VU¹; **Phuong Anh HOANG^{1 2 3}**; Thi Thuan PHAM¹;
Khanh Linh BUI^{1 2}; Thi Ngoc Mai TRAN^{2 3}; Quang Hung LUONG⁴; Thi Hoa Huyen NGUYEN¹

¹College of Health Sciences, VinUniversity, Hanoi, Vietnam;

²Faculty of Nursing and Midwifery, Hanoi Medical University, Hanoi, Vietnam;

³Queensland University of Technology, QLD, 4059, Australia;

⁴The University of Queensland, QLD, 4072 Australia

Background

The prevalence of postpartum depression (PPD) worldwide was approximately 17.22% while the prevalence of postpartum depression among new mothers from 1 to 12 months ranged from 8.2% to 37.1% in Vietnam.

Among Vietnamese women, the risk factors for PPD include personal factors such as level of education, satisfaction about life after birth; family factors such as lack of family support, suffering from sexual violence; and environmental factors such as stressful life events, rural living area or confinement practices. However, there is no evidence of prevalence and predictors of PPD among mountainous mothers.

Objective

This study aims to identify the prevalence of PPD and the risk factors contributing to PPD among women after giving birth in a mountainous area of Vietnam.

Method



A cross-sectional study using convenient sampling method was carried out among women after giving birth in Son La, a province in the northwest region of Vietnam with a diversity of ethnic minorities.

Women who (1) are at least 18-year-old, (2) have delivered their babies within one year, and (3) agree to participate in this study were invited to participate. In total, there were 194 qualified responses from eligible women.

The self-designed questionnaire survey is divided into four parts:

(I) The general *demographic* information of participants

(II) *Perception of PPD* including 15 statements about women's perceptions of PPD (Cronbach's alpha was 0.769)

(III) *Practice for prevention of PPD* consists of 9 questions about the frequency of activities prevent PPD (Cronbach's alpha was 0.825)

(IV) *Edinburgh Postnatal Depression Scale (EPDS)* as the screening tool for PPD which was translated and validated in Vietnamese.

Spearman's correlation analyses were conducted to identify the relations between continuous variables due to the lack of normality of the distribution. A linear regression model was used to determine the relationship between postpartum depression among women after giving birth with other potential variables.

Results

Table 1. Characteristics of participants

Variables	Categories	Frequency	Percentage
Age (Mean SD)		27.2 ± 5.9	
Ethnic groups	Minority	170	87.6%
	Kinh	24	12.4%
Living area	Urban	56	28.9%
	Rural	138	71.1%
Highest level of education	Illiterate	11	5.7%
	Elementary	3	1.5%
	Secondary	67	34.5%
	High school	54	27.8%
	Higher education	59	30.4%
Employment	Unemployment	21	10.8%
	Unskilled labour	131	67.5%
	Skilled labour	42	21.6%
	Single	1	0.5%
Marital status	Divorced/Widowed	1	0.5%
	Married/Living with partner	192	99.0%
Monthly household income per capita	≤ 2,250,000 VND	109	56.1
	> 2,250,000 VND	85	43.9
Health insurance	Yes	17	8.8
	No	177	91.2
Alcohol drinking	Yes	26	13.4
	No	168	86.6
Periodic health check	Less than three times	56	28.9
	At least three times	138	71.1

Figure 2. Prevalence of postpartum depression among women after giving birth in a mountainous area in Vietnam






Table 2. The multiple regression analysis for variables predicting postpartum depression among mountainous women after giving birth

Model	Variable	B	SE	Beta	95% CI	t	p
1	Living area	2.518	0.994	0.193	0.56; 4.58	2.533	0.012
	Monthly household income	-3.477	0.899	-0.292	-5.25; -1.70	-3.868	<0.01
	Periodic Health Examination	-2.746	0.827	-0.211	-4.38; -1.15	-3.32	0.001
2	Living area	2.518	0.906	0.193	0.73; 4.31	2.779	0.006
	Monthly household income	-2.401	0.83	-0.202	-4.04; -0.76	-2.892	0.004
	Periodic Health Examination	-2.331	0.763	-0.179	-3.84; -0.83	-3.053	0.003
	Drinking alcohol	-2.666	1.014	-0.154	-4.67; -0.66	-2.628	0.009
	Perceived infant excessively crying	-2.801	1.098	-0.147	-4.97; -0.64	-2.552	0.012
	Practice	-0.153	0.069	-0.135	-0.29; -0.02	-2.223	0.027
	Perception	-0.200	0.054	-0.23	-0.31; -0.09	-3.68	<0.01
Adjusted R ²		0.383					

Figure 1. Prevalence of postpartum common mental disorders against postnatal screening time points reported by published articles in Vietnam from 2010 to 2020



Conclusion

A large-scale quantitative research should be implemented to establish the association with the potential risk factors. Future qualitative or mix-method studies may provide profound insights into the ***barriers and the mental health demands*** among new mothers in remote locations.

Prospective interventions should aim to ***raise the awareness of mountainous mothers*** about postpartum depression, as well as coping strategies while they have depressive symptoms.