

Feasibility, efficacy and cost of a long-term community-based psychiatric follow-up for People Who Inject Drugs (PWID): the DRIVE-Mind project in Vietnam

Sao Mai LE¹, Philippe TROUILLER², Giang HOANG THI¹, Huong DUONG THI¹, Quang Duc NGUYEN¹, Didier LAUREILLARD^{3,4}, Don DES JARLAIS⁵, Tuyet Thanh NHAM THI⁶, Oanh KHUAT THI HAI⁶, Delphine RAPOUD³, Jean Pierre MOLES³, Nicolas NAGOT³, Laurent MICHEL²

¹ Hai Phong University of Medicine and Pharmacy, 72A Nguyễn Bình Khiêm, Đằng Giang, Ngô Quyền, Hai Phong, Vietnam
² CESP Inserm UMRS 1018, Paris Saclay University, Pierre Nicole Center, French Red Cross, 27 rue Pierre Nicole, 75005 Paris, France
³ Pathogenesis and control of chronic and emerging infections, University of Montpellier, Inserm, Etablissement Français du Sang, University of Antilles, 60 Rue de Navacelles, 34394 Montpellier, France
⁴ Infectious Diseases Department, Caremeau University Hospital, Place du Professeur Robert Debré, 30029 Nîmes, France
⁵ New York University, College of Global Public Health, 665 Broadway Suite 800 NY 10013 New York, USA
⁶ Supporting Community Development Initiatives, 240 Mai Anh Tuan, Thanh Cong Ward, Ba Dinh District, Hanoi, Vietnam

Background

- Adding a mental health problem to drug use increases stigma, severity of both disorders and barriers to accessing care, particularly in places where psychiatric resources are scarce.
- Innovative and alternative interventions are necessary.

→ *The objective of this study was to show that a sustained community-based psychiatric intervention is feasible, efficient in terms of mental health, reduction of drug use, risk behaviors and quality of life, and assess its cost.*

Methods

A total of 233 PWID suffering from mental health problems were proposed a community-based psychiatric follow-up in the city of Haiphong, Vietnam, over 4 years:

- The intervention took place in community-based organization offices with full support of trained peers
- Peers were involved in **research conception, psychiatric screening, questionnaire submission, linkage to care, follow-up and tracing, harm reduction intervention, information & counseling**
- Psychiatric consultations and the treatments, given on sites, were free
- Data were collected every 6 months except during the pike of the COVID epidemic (May 2021 to Mars 2022).
- A control group of 378 PWID free of psychiatric disorders was included in the study during the last year of follow-up to compare severity of drug use and risk behaviors





Main outcomes

- Retention rate
- Psychiatric diagnosis (among **MINI** major depressive episode, psychotic disorder, suicidal risk)
- Psychiatric clinical improvement (**Clinical Global Impression** scale)
- Severity of substance use (heroin injection, regular methamphetamine use, hazardous alcohol use – **audit-C**) (**non-inferiority with control group at M48**)
- Composite risk behavior score (*methadone and injection, syringe/needle sharing, front/backloading, solution for injection sharing, use of equipment to stop bleeding, sharing pipe to smoke methamphetamines, inconsistent condom use with risky primary partner or casual partner*) : **non-inferiority analysis with control group at M48**
- Quality of life (**EQ5D5L**) **non-inferiority analysis with control group at M48**
- HIV/HCV seroconversion compared with control group
- Cost of the intervention



Results

	March 2019		March 2020		COVID Crisis			March 2022		March 2023
Months of Follow-up	M0	M6	M12	M18	M24	M30		M36	M42	M48
Sample size	233	197	170	165	141	//		129	116	111
Retention rate		85%	73%	71%	61%	//		55%	50%	48%
Control group (sample size)								378	→	



SEVERITY of DRUG USE: psychiatric cohort non-inferior to controls in terms of heroin injection, regular methamphetamine use and hazardous alcohol use at M48

HIV/HCV seroconversion: no HIV seroconversion, comparable rate of seroconversion for HCV with controls

COST of the intervention: 78 \$/participant/year

• **CONCLUSION : Sustained community-based psychiatric intervention is extremely efficient, costless and should be widely scaled up, particularly for key populations more often affected by stigma and low income. Nevertheless, access to methadone treatment remains constantly lower among PWID with psychiatric comorbidity compared to controls**