

The Prevalence of Fentanyl in New York City's Unregulated Drug Supply as Measured Through Drug Checking Offered at Syringe Service Programs

Yarelix Estrada¹; Jeffery Sauer¹; Leonardo Dominguez-Gomez¹; Izza Zaidi¹; Andrew J. Trinidad¹; Hannah Helmy¹; Alex Harocopos¹

¹New York City Department of Health and Mental Hygiene, Bureau of Alcohol and Drug Use Prevention, Care, and Treatment, Queens, NY

Background

- In New York City (NYC), the rate of drug poisoning (overdose) deaths has more than quadrupled from 10.2 per 100,000 residents in 2000 to 44.0 per 100,00 residents in 2023.
- Fentanyl has become increasingly involved in overdose deaths, with the percentage of overdose deaths involving fentanyl increasing from 2% in 2012 to 80% in 2020.
- In response, NYC's Department of Health and Mental Hygiene (DOHMH) has strengthened its harm reduction strategies as part of the city's comprehensive overdose response plan.
- To provide real-time information to people who use drugs, DOHMH partnered with community-based syringe service programs (SSP) to conduct a pilot study of point-of-care drug checking.

Methods

- Between November 2021 and December 2023, the pilot study offered drug checking and harm reduction education to participants at community-based SSP operating at fixed sites.
- A drug checking technician was responsible for using the drug checking technology and administering a survey to collect data about the participant's experience.
- Point-of-care drug checking included use of immunoassay test strips and a Fourier-transform infrared (FTIR) spectrometer.




Figure 1. Tools, materials, and equipment used for point-of-care drug-checking.

Source: NYC Health Department's drug checking team (2024)

- While point-of-care provides immediate results to participants, a crucial component of any drug-checking service is the capacity to perform secondary laboratory testing, especially as technicians develop proficiency with drug-checking technologies.
- DOHMH contracted with the Center for Forensic Research and Education (CFSRE) to provide secondary laboratory analysis of collected drug samples, including gas chromatography-mass spectrometry (GC/MS) and liquid chromatography quadrupole time of flight mass spectrometry (LC/QTOF-MS).




Figure 2. Packaged sample to be sent to a secondary testing partner.

Source: NYC health Department's drug checking team (2024)

- Fentanyl positivity was calculated for each drug checking technology (i.e., immunoassay test strip, FTIR, and secondary laboratory testing).
- Fentanyl positivity was further stratified by descriptions of what a sample was sold as, specifically opioids, cocaine, methamphetamine, and benzodiazepines.

Results

- Across the study period, N=397 unique SSP participants brought in n=1083 drug samples to be checked. An additional n=561 drop-off samples were received for a total analytic sample of N=1644.



- Fentanyl positive among opioids samples was very high, although the positivity varied depending on the drug checking technology (FTS: 95.3%; FTIR: 84.7%; secondary laboratory testing: 89.3%).
- None of the cocaine (n=131), methamphetamine (n=29), or benzodiazepine (n=29) samples sent for secondary testing were found to contain fentanyl.

Table 1. Fentanyl-positivity between Fentanyl Test Strip (FTS) and Fourier transform infrared (FTIR) spectrometry

Drug category	Sample Counts		Fentanyl Test Strip Results				
	Sample sold as	Sample confirmed to be	Missing	Negative	Positive	Total	% Fentanyl Positive ^a
	N	N	N	N	N	N	%
Opioid	908	908	14	42	852	894	95.3%
Cocaine	314	309	6	294	9 ^b	303	3.0%
Methamphetamine	62	62	3	59	0	59	0%
Benzodiazepines	59	59	0	58	1 ^c	59	1.7%

Drug category	Sample Counts		Technician-operated FTIR Results ^d				
	Sample sold as	Sample confirmed to be	Missing	No Fentanyl Identified	Fentanyl Identified	Total	% Fentanyl Positive ^a
	N	N	N	N	N	N	%
Opioid	908	908	0	139	769	908	84.7%
Cocaine	314	309	0	308	1 ^e	309	0.3%
Methamphetamine	62	62	0	62	0	62	0%
Benzodiazepines	59	59	0	59	0	59	0%

a: % Fentanyl Positive calculated as: Positive / Total, where Total is the sum of positive (or identified) and negative (or not identified) results.
b: Of these 9 samples, 8 have secondary laboratory results that did not identify fentanyl. We consider these 8 observations to be false positives.
c: Secondary laboratory results for this sample did not identify fentanyl. We consider this observation to be a false positive.
d: FTIR has a limit of detection. Any compound present in the FTIR testing sample in a quantity below the limit of detection cannot be identified.
e: This sample was analyzed during a technician's training phase. The secondary laboratory results for this sample did not identify fentanyl. We consider this observation to be a false positive.

Results (continued)

- A small number of observations (n=5) were omitted from the fentanyl calculations as they represented complex real-world scenarios.
- Each scenario speaks to the critical important of understanding an individual's substance use and how that context can impact drug checking.
- Conversations between drug checking technicians and participants can help reveal critical information that can inform drug checking results.

Table 2. Contextual information revealed through point-of-care drug checking

Participant brought in a sample that was **sold as cocaine** and **stored it in a dope bag**. Drug-checking results indicated the presence of **both cocaine and fentanyl** in the participant's sample.

SSP outreach staff brought in a sample on behalf of a participant. The sample was reported to be **sold as cocaine**. Visual inspection showed that the sample was a **beige powder**. Drug-checking technician and secondary laboratory testing confirmed that it was a dope sample and did not contain cocaine.

Participant brought in a sample that they **thought was cocaine**. Drug checking by technician **identified components that were more consistent with a bag of dope**. After talking with technician, **participant realized** that they had brought in their dope bag instead of their cocaine bag.

Participant brought in samples that were **sold to them as cocaine**. Visual inspection showed that the samples had a purple color and sticky, tar-like texture. The Drug-checking technician and secondary laboratory testing found components consistent with dope samples and did not contain cocaine.

Limitations

- All drug checking samples were voluntarily provided by participants who engaged with established syringe service programs in NYC and thus may not represent the entire unregulated drug supply of NYC.
- A small number of drug samples were missing testing results, most commonly secondary laboratory results.

Conclusions

- A local health department, working in close collaboration with community partners, was able to launch a point-of-care drug checking pilot study.
- The overwhelming majority of samples sold as opioids contained fentanyl.
- The prevalence of fentanyl in samples sold as stimulants was very low.
- Talking to the people who provided the drug checking samples was essential in interpreting findings and informing relevant harm reduction strategies.
- Due to its success, the pilot study will continue as a public health program.

Acknowledgements and Funding

This project was supported by the Centers for Disease Control and Prevention (CDC) of the U.S. Department of Health and Human Services (HHS) as part of Overdose Data to Action (CDC-RFA-CE19-1904) and Overdose Data to Action: LOCAL (CDC-RFA-CE-23-0003). The contents are those of the author(s) and do not necessarily represent the official views of, nor an endorsement by, CDC/HHS or the U.S. Government.

Contact information:

Yarelix Estrada, yestrada1@health.nyc.gov