

INCREASE OF SYNTHETIC OPIOID USE: TWO EVIDENCE-BASED AND LOW TRESHOLD INTERVENTIONS TO PREVENT DRUG RELATED DEATHS

BACKGROUND

Synthetic opioids are a major contributor to drug-related deaths in different countries (2022, USA 68% of 107.941 drug-related deaths¹; 2023, Canada 82% of 8.049 opioid-related deaths²). The current European drug report shows that 20 countries in Europe are already confronted with synthetic opioids such as Nitazene and Fentanyl. Germany saw 2023 the highest number of drug-related deaths ever (2.227 cases). Synthetic opioids was registered in 76 cases³. **Two german studies** (Take Home Naloxone⁴ and Rapid fentanyl testing⁵) **have proven measures for preventing opioid-related drug emergencies and drug deaths.**

DESCRIPTION

- **RAFT** (Rapid fentanyl Testing): The multi-centred study was carried out in 17 drug consumption rooms. With easy-to-use rapid tests, 1.405 heroin samples were tested. Users receive the result quickly, anonymously and have no loss of heroin. If the test is positive, fentanyl-related counselling is provided. **The main goal was to raise awareness for fentanyl and support fact-based consumption decisions.**
- **NALtrain** (Take Home Naloxone): It is a nationwide program that trains opioid users and employees in the **use of naloxone** and provides them with the medication in case of an emergency.

LESSONS

- **RAFT** Within 6 months, **3% of the samples tested positive for fentanyl**. After fentanyl-related counseling, one in five users took advantage of harm reduction measures such as dose splitting.



Naloxone
nasal
spray



- **NALtrain** Around 870 people were trained as multipliers in 74 training sessions by the end of 2023. Around 2.208 participants were provided with THN. **110 uses of naloxone** were documented. No deaths were recorded.

CONCLUSION

Both projects shows that there is a great interest and willingness among employees and users to implement new, low-threshold and effective harm reduction measures.

BOTH INTERVENTIONS ARE HIGHLY EFFECTIVE IN PREVENTING DRUG EMERGENCIES AND DRUG-RELATED DEATHS.

In view of the fact that drug checking cannot be implemented politically in many countries, **fentanyl rapid tests are one effective way of reducing harm**. **Take-home naloxone is an inexpensive life-saving medication** that every user should be equipped with. Both projects are building blocks of effective and evidence based strategies that can be applied regardless of the economic status of the countries.

¹ NIDA, Online 14-07-2024

² Government of Canada, Online 14-07-2024

³ Bundesdrogenbeauftragter, Online 14-07-2024

⁴ further information: www.naloxonetraining.de *)

⁵ further information: www.drogenkonsumraum.de/raft *)

*) in continous processing