

How to improve access to methadone treatment among people with co-occurring psychiatric disorder and opioid use? Answers from a cross sectional survey conducted in Haiphong, Vietnam.

Philippe Trouiller¹, Sao Mai Le², Giang Hoang Thi², Huong Duong Thi², Oanh Khuat Thi Hai³, Khue Pham Minh², Roselyne Vallo⁴, Catherine Quillet⁴, Thuy Linh Nguyen², Quang Duc Nguyen², Tuyet Thanh NhamThi³, Jonathan Feelemyer⁵, Vinh Vu Hai⁶, Jean-Pierre Moles⁴, Didier Laureillard^{4,7}, Don C. Des Jarlais⁵, Nicolas Nagot⁴, Laurent Michel¹ and Delphine Rapoud⁴ for the DRIVE Study Team

¹ CESP Inserm UMRS 1018, Paris Saclay University, Pierre Nicole Center, Fench Red Cross, 27 rue Pierre Nicole, 75005 Paris, France

² Hai Phong University of Medicine and Pharmacy, 72A Nguyễn Bình Khiêm, Đằng Giang, Ngô Quyền, Hai Phong, Vietnam

³ Supporting Community Development Initiatives, 240 Mai Anh Tuan, Thanh Cong Ward, Ba Dinh District, Hanoi, Vietnam

⁴ Pathogenesis and control of chronic infections, Inserm, Etablissement Français du Sang, University of Montpellier, 60 Rue de Navacelles, 34394 Montpellier, France

⁵ New York University College of Global Public Health, 665 Broadway Suite 800
NY 10013 New York, USA

⁶ Dept of Infectious and tropical diseases, Viet Tiep Hospital, Số 1 Đường nhà thương - Quận Lê Chân, Haiphong, Vietnam

⁷ Infectious Diseases Department, Caremeau University Hospital, Place du Professeur Robert Debré, 30029 Nîmes, France

Backgroud. People with co-occurring psychiatric disorder and opioid use take particularly advantage from methadone treatment but seem to have less access to it than other populations. To improve this situation, we need to quantify the scale of the problem and identify the main obstacles to access to methadone. In this study, we aimed to explore the relationship between psychiatric disorders and access to methadone treatment in a cohort of people who inject drugs (PWID) in Haiphong, Vietnam.

Methods. A cross-sectional survey was conducted in 2020 among a cohort of PWID recruited in the city of Haiphong by respondent-driven sampling. Data on socio-demographic characteristics, drug use, quality of life, and access to treatment were collected using face-to-face questionnaires by trained interviewers. PWID were screened by trained psychiatrists for depression, psychotic disorder and suicidality, using the MINI questionnaire.

Results. Among the 492 PWID enrolled in this study, 18% were not taking methadone despite their intention to do so. Mean age was 45 years and 93% were male. People with psychiatric disorder were almost half as likely to be treated with methadone (Odds Ratio: 0.54, IC95% [0.34 – 0.88]). This relationship was mostly mediated by two factors in our multivariate analysis: reporting being a woman or a transgender person, or reporting poor mobility. On the other hand, income, age and heroin use did not statistically mediated this relationship.

Conclusions. New strategies to improve methadone coverage among PWID with dual disorders should include take home methadone (so far only daily delivery is available in Vietnam), peer support, and easier access by women and transgender people to the methadone clinics.

Key words: psychiatric disorder, methadone, access to care