

Feedback on staff's readiness for implementation

M. HARTVEIT^{1,2}, H. NORDIN³, T. RUUD⁴

1 Fonna Hospital Trust, Haugesund, Norway. 2 University of Bergen, Bergen, Norway. 3 Stavanger Hospital Trust, Stavanger, Norway
4 University of Oslo, Oslo, Norway

FINAL ID

INTRODUCTION

Care providers constantly assess helping and hindering conditions during implementation of evidence-based practices [1].

Implementation leadership imply facilitation and support tailored to the care providers' changing needs [2].

However, the empirical grounding for these assumptions are week.

AIM

- We aimed to
- a) test if staff's reported readiness is associated with implementation outcomes at different phases in the implementation, and
 - b) explore the usefulness of feedback on such readiness for implementation leadership.

METHOD

Cluster-RCT including 39 Norwegian mental health care centers implementing evidence-based practices for psychosis treatment.

Intervention: Training, implementation support and feedback every sixth month on staff's readiness and implementation outcomes.

Data: Implementation Process Assessment Tool (IPAT) questionnaire [3] and implementation outcomes by fidelity scales.

Structured individual interviews with leaders post-hoc.

The Implementation Process Assessment Tool (IPAT):




Figure 1: Illustration of the theoretical grounding for the Implementation Process Assessment Tool (IPAT) [3]. Factors of the main CIR domains Outer setting, Inner setting, Intervention characteristics and Characteristics of individuals [1] are constantly being interpreted by care providers, implying readiness at two levels; the individual and the collective. The degree of readiness and the progression of stages of change are expected to interact.

Examples from the 27 items IPAT [3] :

Item 4: *I have discussed with colleagues how this new practice will work in our unit.*


Item 16: *I believe the patients will benefit from the improvement.*

Item 26: *I feel confident that we [your team] have the necessary knowledge and experience of systematic improvement.*

RESULTS

Staff's readiness was positively affected by the implementation supportive intervention (t(140)=2.22, p=0.028) [5].

Staff's reported readiness (IPAT-score) associated with implementation outcome (fidelity-score) at 12 months: (B(CI)=0.4 (0.0, 0.7), p=0.045) and 18 months: (B(CI)=0.7 (0.2, 1.1), p=0.004) [4].



Leaders' experiences of usefulness and challenges using the IPAT feedback (N=37):

Helps to map the situation and staff's needs

(The IPAT feedback) did not led to decisions but raised our awareness. It revealed lack of engagement in the out-patient clinic. We managed to involve them.

(In retrospect) I think IPAT-feedback would have been useful. Have experienced great frustration among the leaders for failing the implementation.

Lack implementation skills and knowledge

I found IPAT-reports useful, but to profit from them I need to understand them better.

I see the usefulness and understand what it means but find it difficult to translate it to action.

The implementation facilitator's competence was useful for interpreting the IPAT-feedback.

Involvement and integration

The information was useful but is somehow not in line with how we do it. A few individuals do this implementation - no collective ownership.

I've discussed the IPAT feedback with my two clinical advisors, but I am not involved in the implementation process.

It is the clinicians that know about this implementation. I know little about it.

Figure 2: Themes and statements from leaders about the usefulness and challenges using the IPAT feedback.




Figure 3: Number of leaders reporting to use the IPAT feedback and how many reported active adaption of the implementation support based on the feedback. «New» leader = leadership change during the study period.

CONCLUSIONS

Larger success in implementation can be achieved by repeated feedback on staff's IPAT-scores. However, the gain of such feedback relies on sufficient implementation leadership knowledge and skills.

Implications:

- We should pay more attention to care providers' readiness throughout the implementation process to complete and sustain the new practice.
- Implementation leadership training should be emphasized.
- Further research on informed implementation leadership is needed.

DECLARATIONS

Ethical approval: The study was approved by The Regional Committee for Medical and Health Research Ethics in Southeastern Norway (Reg.no. REK 2015/2169) and the Data protection officer for each participating hospital trust.

Registration: [ClinicalTrials.gov](https://clinicaltrials.gov/ct2/show/study/NCT03271242) Identifier: [NCT03271242](https://clinicaltrials.gov/ct2/show/study/NCT03271242)

Funding: The study was funded by the South-Eastern Regional Health Authority (Helse Sør-Øst HF) in Norway (Grant No. 2015106).

Declaration of interests: None.

REFERENCES

1. Damschroder L et al. "The updated Consolidated Framework for Implementation Research based on user feedback." *Implementation science* 17.1 (2022): 75.
2. Nordin HMA et al. "Implementing Clinical Guidelines for the Treatment of Psychosis: The Frontline Leaders' Point of View. A Qualitative Study." *Journal of Healthcare Leadership* (2024): 93-104.
3. Hartveit M et al. "Measuring implementation: development of the implementation process assessment tool (IPAT)." *BMC Health Services Research* 19 (2019): 1-12.
4. Hartveit M et al. "Can systematic implementation support improve programme fidelity by improving care providers' perceptions of implementation factors? A cluster randomized trial." *BMC Health Services Research* 22.1 (2022): 808.
5. Ruud T et al. "The effect of intensive implementation support on fidelity for four evidence-based psychosis treatments: a cluster randomized trial." *Administration and Policy in Mental Health and Mental Health Services Research* 48 (2021): 909-920.



CONTACT INFORMATION

Corresponding author:

Miriam Hartveit (Miriam.Hartveit@helse-fonna.no)

Valen hospital, Helse Fonna HF

N-5450 Valen

Norway