

A case of hyperreactio lutenalis complicated by mirror syndrome presenting with characteristic ultrasound findings

2. School of Health Sciences Department of Family and Child Nursing, and Midwifery, Shinshu University, Matsumoto, Nagano Prefecture, Japan



[Introduction]Hyperreactio lutealis (HL) is a benign condition characterized by bilateral, multicystic enlargement of the ovaries due to hCG stimulation during pregnancy. We report a case of HL associated with Mirror Syndrome, which is characterized by fetal hydrops, maternal edema, and placental edema. [Case Report] A 29-year-old woman, gravida 2 para 0, became pregnant through frozen-thawed embryo transfer. At 12 weeks of pregnancy, cystic hygroma was detected by her previous physician. At 15 weeks of pregnancy, both of her ovaries were found to be enlarged to 9 cm, and her serum hCG level was elevated to more than 1,000,000 mIU/mL. Although malignancy was ruled out by MRI, further evaluation and treatment were recommended, and she was transferred to our hospital at 16 weeks of pregnancy. Ultrasound examination revealed fetal hydrops, and the placenta appeared edematous and thickened. Both ovaries were enlarged to 10 cm with a multicystic structure, showing a "spoke-wheel appearance," leading to a diagnosis of HL. The mother showed elevated blood pressure and generalized edema, and blood tests revealed hemoglobin at 8.2 g/dL, hematocrit at 25.1%, indicating hemodilution, and markedly elevated serum hCG at 1,322,000 mIU/mL, raising suspicion of Mirror syndrome. Two days after her transfer, intrauterine fetal death occurred, and the fetus was delivered the following day, but the placenta did not expel naturally. Due to worsening maternal hypertension and edema, as well as the deterioration of her general condition, a uterine evacuation was performed two days later to remove the placenta. After the procedure, her condition stabilized, and her serum hCG level dropped to 17,047 mIU/mL on the 8th postoperative day, allowing for discharge. Her ovarian condition is being monitored on an outpatient basis. [Conclusion] HL can be misinterpreted as an ovarian tumor requiring surgical removal due to its rapid enlargement during pregnancy, but if diagnosed based on typical clinical course and imaging, conservative management is recommended as it spontaneously regresses. It is important to note that, as in this case, hyperhCGemia associated with Mirror syndrome can also be a trigger for the development of HL.

Hemodilution, Abnormally High hCG, Normal tumor maker

Name of the Authors: Ryoichi Asaka
Affiliation: Department of Obstetrics and Gynecology, Shinshu University
I have no financial relationships to disclose.