

OBJECTIVE

- Selective fetal growth restriction (sFGR) complicates 10-15% of monochorionic (MC) twin gestations and optimal management remains unknown.
- Umbilical artery (UA) waveforms in sFGR fetuses are used to determine types (Type 1- normal, Type 2- persistently abnormal, Type 3- intermittently abnormal).
- We aimed to uncover outcomes of a standardized protocol for expectant management in sFGR with abnormal UA Doppler waveforms (types 2 and 3).

METHODS

- A retrospective cohort analysis of MC twin pregnancies with sFGR managed and delivered at a single tertiary center (2015-2023).
- Excluded: patients undergoing laser. All were admitted at 26-28 weeks, received antenatal steroids, and underwent intensive monitoring (twice weekly ultrasounds, NST q8h).
- Clinical data were obtained from EMR.

Table 1.

Outcome	Type 2 sFGR (n=13)	Type 3 sFGR (n=20)	P-value
Birthweight (g)	921 ± 317	1213 ± 479	0.09
Birthweight (%tile)	12.2 ± 8.0	14.5 ± 15.0	0.71
1 min Apgar	5 [3, 8]	7 [5, 8]	0.14
5 min Apgar	7 [7, 8]	8 [8, 9]	0.09
Arterial cord pH	7.29 ± 0.07	7.35 ± 0.05	0.09
IVH (%)	15	0	0.16
RDS (%)	46	65	0.29
NEC (%)	8	5	1.0
ROP (%)	8	5	1.0
Neonatal death (%)	8	0	0.41
NICU LOS (days)	108 ± 49	72 ± 74	0.11

RESULTS

- 33 pregnancies with sFGR and abnormal UA Dopplers were managed. 13 (39%) were Type 2, 20 (61%) were Type 3.
- No difference in GA at diagnosis. GA at delivery was higher in Type 3 (31 [29,33] vs 29 [27,31] weeks, *p*=0.03).
- Non-reassuring fetal status was the most common indication for delivery in both groups (77% vs 65%).
- Neonatal outcomes were similar. One neonatal death occurred in Type 2.

CONCLUSIONS

- In this cohort, Type 3 sFGR was associated with later gestational ages, but delivery for non-reassuring fetal status occurred in ~2/3 of patients. Neonatal survival >98%.
- Standardized inpatient protocol may be associated with favorable perinatal outcomes.