


Inter-rater Reliability of Sonologists Assessing MUSA Uterine Features in Symptomatic Women at a Tertiary Hospital in the Philippines

Presenter: Massomeh VC. Shamsoddin - Padiernos, MD., FPOGS
Co-Author: Cherry Go - Palomar, MD., FPOGS, FPSUOG

- Research Context:**
- ✓ Clinical and Ultrasonographic Features of Adenomyosis
 - ✓ MUSA Framework Reproducibility as a Standardized Diagnostic Tool
 - ✓ Sonologists Assessing MUSA Features
 - ✓ Analytical Cross-Sectional Study Design
 - ✓ 22 Symptomatic Women Aged 25-58

Parameter	% Agreement	Gwet's AC	95% Confidence Interval	Interpretation
Interrupted Junctional Zone (JZ)	95.45%	0.95	0.90-1.00	Almost Perfect
Overall Diagnosis	87.27%	0.85	0.73-0.97	Strong Consensus



MUSA Criteria Unlock Strong Concordance for Adenomyosis Diagnosis at AC = 0.85

CONCLUSION

INTERRUPTED JUNCTIONAL ZONE (JZ)

The most dependable marker

EXPERT AGREEMENT IS HIGH

Overall consensus is 87.27%

GWET'S AC UNBIASED MEASURE

Use Gwet's AC (not Fleiss' Kappa) when feature prevalence is skewed

