

# Mass Spectrometry-Based Proteomics of Polymethyl Methacrylate-Embedded Bone

Annika Topitsch<sup>1,2,3</sup>, Tim Halstenbach<sup>1,4</sup>, Annekathrin Kopp<sup>5</sup>, Annette Lindner<sup>4</sup>, Susanne Nahles<sup>5</sup>, Oliver Schilling<sup>1</sup>, Tobias Fretwurst<sup>4,6</sup> and Katja Nelson<sup>4</sup>

<sup>1</sup>Institute for Surgical Pathology, University Medical Center Freiburg, Germany


<sup>2</sup>Spemann Graduate School of Biology and Medicine (SGBM), University of Freiburg, Germany

<sup>3</sup>Faculty of Biology, University of Freiburg, Germany

<sup>4</sup>Department of Oral and Maxillofacial Surgery, University Medical Center Freiburg, Germany

<sup>5</sup>Department of Oral and Maxillofacial Surgery, Charité Berlin, Germany

<sup>6</sup>Department of Oral Medicine, Infection and Immunity, Harvard School of Dental Medicine, Boston, USA



## Introduction

Polymethyl methacrylate (PMMA) embedding media are widely used for the preservation of hard tissues such as bone and teeth. Various studies have shown the applicability of PMMA-embedded specimens for histological and immunohistochemical investigations [1-3]. However, for further research on the cellular and molecular mechanisms in bone, in-depth and high-throughput analysis techniques are essential. One such technique is mass spectrometry (MS)-based proteomics, which is used to identify and comparatively quantify the proteins in a biological sample. At present, clinical proteomics largely focuses on soft tissues or body fluids, while the use of bone as a sample is not well established [4-6]. This study is the first to answer the question: **Are PMMA-embedded bone specimens suitable for MS-based proteomics?**

## Methods

**Samples:** Porcine jawbone cut into 5 mm cubes

**Embedding:** Technovit 9100 or fresh-frozen in liquid nitrogen


**Treatment:** Decalcification in HCl for 17 days for half of the samples

**Protein extraction:** Grinding using Precellys MK28 beads; sonication; incubation at 95°C in SDS

**Digestion:** Reductive alkylation using TCEP and CAA; digestion with trypsin protease

**Labeling:** Chemical labeling using isobaric tandem mass tags (TMTs)


**Measurement:** Data-dependent acquisition on a Q Exactive Plus mass spectrometer



## Results

### 1 Quantification of Proteins


- Quantified proteins cover a wide range of abundances in all samples and replicates
- Homogenous distribution of intensities in TMT labeling channels



Left: Identified proteins ranked by logarithmic abundance. Right: Distribution of logarithmic abundances per TMT channel after median normalization. Data analysis was performed using the FragPipe software.

### 2 Identification of Proteins


- 1096 proteins identified across all samples  
→ similar numbers for Technovit and fresh-frozen samples
- Three most highly abundant proteins:  
Collagen type 1 alpha 1 and 2, collagen type 2 alpha 1



Left: Numbers of identified proteins per sample. Right: Percentage of the three most highly abundant proteins per sample.

### 3 Proteome Comparisons


- Clustering in PC analysis → Considerable differences in proteome of Technovit and fresh-frozen samples
- High reproducibility and comparability within workflows
- Numerous significantly differentially abundant proteins



Left: Principal component (PC) analysis showing the clustering of workflow replicates. Right: Differential abundance analysis comparing the effects of embedding techniques and decalcification reveals substantial fold changes (FCs) of proteins that are significantly differentially abundant between workflows. Statistical analysis was performed in R using the limma package.

### 4 Effect of Decalcification

- Varying numbers of identified proteins within Technovit replicates
- Clustering in PC analysis  
→ Considerable differences between samples
- Additional proteins identified




Left: Numbers of identified proteins per sample. Middle: PC analysis showing the clustering of workflow replicates. Right: Overlap of identified proteins of Technovit-embedded (top) and fresh-frozen (bottom) samples for all sample preparation workflows. Data analysis was performed using the MaxQuant software.

## Conclusion

- High numbers of identified proteins, independent of embedding technique and sample treatment
- Both, embedding technique and sample treatment, affect proteome composition
- Consistent sample handling ensures high reproducibility and comparability
- Decalcification can increase the numbers of identified proteins, but is not recommended for Technovit-embedded samples  
→ **Yes, MS-based proteomics can be performed on PMMA-embedded bone specimens**

## Outlook

Laser microdissection of Technovit sections



Microscopic isolation of bone tissue from Technovit sections  
→ Challenging task to detect low-abundant proteins

## References

- Willbold, E., Reedmann, M., Jeffreis, R. & Witte, F. Histology and research at the hard tissue-implant interface using Technovit 9100 New embedding technique. *Acta Biomater.* 6, 4447-4455 (2010).
- Wittenburg, G., Volkel, C., Mai, R. & Lauer, G. Immunohistochemical comparison of differentiation markers on paraffin and plastic embedded human bone samples. *J. Physiol. Pharmacol.* 60, 43-49 (2009).
- Yang, R., Davies, C. M., Archer, C. W. & Richards, R. G. Immunohistochemistry of matrix markers in Technovit 9100 New-embedded undecalcified bone sections. *Eur. Cells Mater.* 6, 57-71 (2003).
- Pauwels, F. & Gevaert, K. Mass spectrometry-based clinical proteomics – a revival. *Expert Rev. Proteom.* 18, 411-414 (2021).
- Halstenbach, T., Nelson, K., Igthaut, G., Schilling, O. & Fretwurst, T. Impact of peri-implantitis on the proteome of crevicular fluid: A pilot study. *J. Periodontol.* 1-13 (2023).
- Fretwurst, T., Fritschler, I., Rothweiler, R., Nahles, S., Altmann, B., Schilling, O. & Nelson, K. Proteomic profiling of human bone from different anatomical sites – A pilot study. *Proteomics Clin. Appl.* 16 (2022).

## Acknowledgements

OS acknowledges funding by the German Research Foundation (DFG), SCHI 871/17-1, NY 90/6-1, SCHI 871/15-1, GR 4553/5-1, PA 2807/3-1, project-ID 431984000 – SFB 1453 (P12), project-ID 441891347 - SFB 1479 (S1), project-ID 423813989 – GRK2606, project-ID 32297 7937 – GRK 2344, the ERA PerMed program (BMBF, 01KU1916, 01KU1915A), and the German Consortium for Translational Cancer Research (project Impro-Rec).

The authors declare no competing financial interest.

Illustrations were created with BioRender.com.

