

Two Cases of Bone Augmentation with Carbonate Apatite

Shunichi Shibazaki, Hidetaka Miyashita, Satoru Morikawa, Taneaki Nakagawa


Department of Dentistry and Oral Surgery, Keio University School of Medicine, Tokyo, Japan

Background

Guided Bone Regeneration is performed to reconstruct the alveolar crest to an appropriate morphology to achieve more functional and aesthetic restoration. Here, we present two cases of GBR with carbonate apatite (Cytrans[®]Granules).

Case1

- 59-year-old Female
- Chief complaint : cannot chew



First visit examination

Case2

- 66-year-old Female
- Chief complaint : needs treatment for periodontitis




First visit examination


Two Cases of Bone Augmentation with Carbonate Apatite

Shunichi Shibazaki, Hidetaka Miyashita, Satoru Morikawa, Taneaki Nakagawa

Case1



Case2



Materials and Methods

Case1

A staged approach GBR was performed for the number #10-14 missing. (a)(b) Clinical appearance of horizontal and vertical ridge defect ; (c)(d) Placement of the mixture of carbonate apatite and autogenous bone with titanium mesh.

Case2

A staged approach GBR was performed for the number #3,4 missing. (e)(f) Clinical appearance of horizontal and vertical ridge defect ; (g)(h) Placement of the mixture of carbonate apatite and autogenous bone with titanium mesh.

Two Cases of Bone Augmentation with Carbonate Apatite





Shunichi Shibazaki, Hidetaka Miyashita, Satoru Morikawa, Taneaki Nakagawa

Results

Case1

About 6 months after GBR, implant placement was planned. Bone augmentation was achieved horizontally in areas #10-12 and vertically in area #14.



Pre-treatment



areas #10-12

area #14

6M after GBR



areas #10-12

area #14

Two Cases of Bone Augmentation with Carbonate Apatite





Shunichi Shibazaki, Hidetaka Miyashita, Satoru Morikawa, Taneaki Nakagawa

Results

Case2

About 6 months after GBR, implant placement was planned. Bone augmentation was achieved vertically in area #3 and horizontally and vertically in area #4.





Pre-treatment



area #3

area #4

6M after GBR



area #3

area #4



Two Cases of Bone Augmentation with Carbonate Apatite

Shunichi Shibazaki, Hidetaka Miyashita, Satoru Morikawa, Taneaki Nakagawa



Case1

Case2

Immediately
after GBR



6M after GBR



Results

When comparing the immediate postoperative period to 6 months later, there was little resorption of the bone substitute in the waiting period.

Conclusion

While primary closure and stability of the graft material are prerequisites for successful GBR, we consider carbonate apatite is a good material based on the results of this study. Although the number of cases is still small, we will continue to report on this material in the future.