

# Immediate implant placement with CBCT surgical planning: A case report

Torres B.,D.\* Soto S., V. \*\*Pimentel H.,J.\*\*\*

March 16-18,2023  
Phoenix, Arizona

## Introduction:

Dental implant placement provides a predictable and safe treatment option for oral rehabilitation. A successful treatment must be based on the final prosthesis, it is recommended the use of a CBCT designed surgical guide and mockup. These elements minimize the risks in terms of the ideal position and the potential damage to anatomical structures, in addition the immediate implant placement reduces the number of interventions thus rehabilitation time, also help us to preserve and maintain the soft tissue.

\* DDS, Advanced Restorative Dentistry program resident, UNAM, México.

\*\* DDS, Periodontics Faculty in Advanced Restorative Dentistry program, UNAM, México.

\*\*\* DDS, Prosthodontics Faculty in Advanced Restorative Dentistry program, UNAM.

# Case Report:

A 49-year-old female patient with periodontal diagnosis of gingivitis induced by dental plaque.


Unfavorable prognosis non-restorative root canal treatment



Fig. 1. Occlusal view with the initial conditions

Seibert II defect in 25-26

Atraumatic extractions + immediate implant placement



Implant placement in healed site

Fig. 2. Treatment plan

# Treatment Sequence:

# CBCT Surgical Planning:





Fig. 3. Diagnostic Wax up and the guide design





Fig. 4. Implant 12 planning position





Fig. 6. Implant 15 y 14 planning position





Fig. 5. Implant 22 planning position





Fig. 7. Implant 24 y 26 planning position

# Treatment Sequence:

## Surgical Phase



Fig. 8. Immediate post extraction




Fig.9. Guide try in




Fig. 10. 6 implant placed



Fig. 11. Implants with healings screws



Fig. 12. Gap filled with particulate bovine-derived xenograft and a collagen matrix




Fig. 13. Wound closure was done with 5-0 polyglycolic acid

## Results:




Fig. 14. Post operative follow up



Fig. 15. Provisionalization




Fig. 16. Soft tissue augmentation with connective tissue graft

## Conclusions:

- ✓ The immediate implant placement helps to optimize rehabilitation time since it reduces the number of surgical interventions.
- ✓ The use of a surgical guide increases success rates by reproducing the ideal position based on the previously planned restoration.

## References:

