



Implant Success Rate and Marginal Bone Loss Following Vertical and Horizontal Guided Bone Regeneration: A Retrospective Analysis



Joseph Kan DDS, MS¹, Samuel Sauza DDS², Joseph Lee DDS¹, Sunee Limmeechokchai DDS, MSD¹, Sasiya Bhumpattarachai DDS, MSD¹, Shi Yin DDS, MS³, Ji Yeon Chung DDS², Aladdin Al-Ardah BDS, MS¹, Jaime Lozada DDS¹

¹Advance Education in Implant Dentistry, School of Dentistry, Loma Linda University, Loma Linda, California, USA

²Department of General Dentistry, School of Dentistry, Loma Linda University, Loma Linda, California, USA

³Private Practice, Seattle, Washington, USA

BACKGROUND

Therapy for partially dentate and edentulous patients are routinely resolved with endosseous implants. The general guidelines and requirements for implant placement are commonly limited to the availability of bone. The increased use of implants resulted in the increase of bone augmentation procedures for hard tissue defects. Guided bone regeneration (GBR) utilizes barriers for adequate revascularization, space maintenance, and stability for graft materials. Although GBR may provide satisfactory volumetric bone regeneration for implant placement, the long-term success of implants in grafted sites remain unclear.

PURPOSE

Question: *Do implants placed in horizontally or vertically guided bone regenerative sites maintain marginal bone level from the time of implant placement?*

The aim of this study is to assess the marginal bone stability, success rate, and survival rate of implants in GBR sites.



MATERIALS & METHODS



A chart review of all patients who received GBR and then implant placement in both at Loma Linda School of Dentistry Center for Implant Dentistry (LLUSD) between 2000 to 2020 was conducted.

INCLUSION CRITERIA

- Age 18 years or older
- Available pre-/post-operative radiographic images
- Radiograph image following implant placement
- Radiograph image following restorative insertion
- Radiograph after at least 6 months of functional loading
- Implants removed at the time of 2nd stage procedure due to failure

DATA COLLECTION VARIABLES

- Demographics
- Date of GBR procedure
- Type of bone graft
- Type of membrane
- Date and radiograph of implant placement
- Date and radiograph of prosthesis loading
- Date of latest radiograph




Fig 1. Pre-op Radiograph




Fig 2. Post-op Radiograph




Fig 3. Sequence of radiographs, calibrated for measurement



MATERIALS & METHODS



Test Group 1 - Vertical GBR

Radiograph pre-/post-op $\geq 3\text{mm}$ gain in vertical direction

Test Group 2 - Horizontal GBR

Radiograph pre-/post-op $\geq 3\text{mm}$ gain in horizontal direction

DATA ANALYSIS

- Radiographic measurements by periodical radiographs :
 - Time of bone graft (T_0)
 - Time of implant placement (T_1)
 - Time of prosthetic load (T_2)
 - Time of latest follow-up (T_3)
- Measurements made on Apple Keynote (Cupertino, CA)

Criteria for implant failure:

- $\geq 30\%$ bone loss at the time of follow-up with/without implant removal
- Implants removal at the time of 2nd stage uncoverly




Fig 4. Keynote Measurement Diagram

Measurement (units in points)

X-axis horizontal lines drawn at the level of implant platform and apex = Total length of implant (z)

X-axis horizontal lines drawn at level of bone at $T_1 = (x)$

X-axis horizontal lines drawn at level of bone at $T_3 = (y)$

Equation for bone level change in percentage:

$$(x-y)/z*100\% = q$$



MATERIALS & METHODS




Fig 5. Cluster bar chart of implant failures over time

	Group 1 (Vertical)	Group 2 (Horizontal)
Number of patients	37	51
Total implants (<i>n</i> = 201)	79	122
Mean Follow-up Time	75.6 mos	67 mos
Mean bone loss (%)	29.4%	9.8%
Implant Removal	12	2
Marginal Bone Loss ≥ 30%	30	7
Overall Failure Rate	37.97%	5.74%

Table 1. Results of implant included in study

	Group 1	Group 2
Ti-Mesh	25	4
Collagen Membrane	20	112
PTFE Membrane	23	5

Table 2. Overall Implants Barrier Types

	Group 1	Group 2
Allograft	29	34
Xenograft	2	14
Autograft + Allograft	14	16
Autograft + Xenograft	7	0
Allograft + Xenograft	11	32
Autograft + Allograft + Xenograft	12	24

Table 3. Overall Implants Bone Graft Types



RESULTS & CONCLUSION



	Group 1 (Vertical)	Group 2 (Horizontal)
Total Failures	30	7
Ti-Mesh	10	—
Collagen Membrane	13	5
PTFE Membrane	7	2

Table 4. Implant Failure based on Barrier Type

	Group 1 (Vertical)	Group 2 (Horizontal)
Allograft	16	—
Xenograft	—	—
Autograft + Allograft	4	2
Autograft + Xenograft	2	—
Allograft + Xenograft	5	1
Autograft + Allograft + Xenograft	3	4

Table 5. Implant Failure based on Bone Graft Type

RESULTS

- 57 Females, 29 Males
- Mean age 60, range age 20-81
- Group 1 Mean follow-up 75.6mos (0-145)
- Group 2 Mean follow-up 67mos (12-168)

CONCLUSIONS

- **Within the confines of this study, about 1/3 of vertical GBR cases experienced significant bone loss around implants over the course of 6 years**
- **Horizontal GBR, on the contrary, remains relatively successful over 5 years**

REFERENCE

1. Nyman S, Lindhe J, Karring T, Rylander H. New attachment following surgical treatment of human periodontal disease. J Clin Perio, 1982.
2. Gottlow J, Nyman S, Lindhe J, Karring T, Wennstrom J. New attachment formation in the human periodontium by guided tissue regeneration. Case Reports. J Clin Perio 1986.
3. Dahlin C, Sennerby L, Lekholm U, Linde A, Nyman S. Generation of new bone around titanium implants using a membrane technique: An experimental study in rabbits. Int J Oral Maxillofac Implants, 1989.
4. Buser D, Bragger U, Lang NP, Nyman S. Regeneration an enlargement of jaw bone using guided tissue regeneration. Clin Oral Implants Res 1990.
5. Urban IA, Jovanovic SA, Lozada JL. Vertical ridge augmentation using guided bone regeneration (GBR) in three clinical scenarios prior to implant placement: A retrospective study of 35 patients 12 to 72 months after loading. Int J Oral Maxillofac Implants, 2009.
6. Urban IA, Nagursky H, Lozada JL, Nagy K. Horizontal ridge augmentation with a collagen membrane and a combination of particulate autogenous bone and inorganic bovine bone derived mineral: A prospective case series in 25 patients. Int J Periodontics Restorative Dent, 2013.
7. Hammerle CHF, Jung RE, Feloutzis A. A systematic review of the survival of implants in bone sites augmented with barrier membranes (guided bone regeneration) in partially edentulous patients. J Clin Periodontol, 2002.
8. Salvi GE, Monje A, Tomasi C. Long-term biological complications of dental implants placed either in pristine or in augmented sites: A systematic review and meta-analysis. Clin Oral Impl Res, 2018.
9. Urban IA, Montero E, Monje A, Sanz-Sanchez I. Effectiveness of vertical ridge augmentation interventions: A systematic review and meta-analysis. J Clin Periodontol, 2019.
10. Chiapasco M, Zaniboni M, Boisco M. Augmentation procedures for the rehabilitation of deficient edentulous ridges with oral implants. Clin Oral Implants Res, 2006