

The effects of bone grafting on immediate implant primary stability in a simulated model

Satyavrat Arya, Leena Garg, Sebastiano Andreana

Buhite-DiMino Center for Implant Dentistry and Advanced Dental Education University at Buffalo School of Dental Medicine

Introduction

Research question: Does bone grafting contributes to the overall success of dental implants immediately placed in tooth extraction sockets?

Aims and Objectives:

- 1) To evaluate the primary stability of dental implants placed in extraction sockets with or without bone graft.
- 2) To assess the effect of different bone graft materials on primary stability.

Hypothesis: Bone grafting affects primary stability of the implants placed immediately in extraction sockets.

Material and methods

In vitro experimentation with a sample size of 20 osteotomies (2.75 * 10mm) using porcine ribs to simulate clinical conditions.

- Root form endosseous Implant (3.8 x 10mm)
- The RFA equipment
- Alloplast putty
- Xenograft particulate
- 6 mm bone Trephine



The extraction socket defects were created modifying a previously published technique. 6mm of bone was removed coronally from porcine ribs simulating the socket defect.⁴





Figure1:

- (A) No socket defect (Control) (Gc);
- (B) Socket defect with no bone graft (Test) (GT0);
- (C) Socket defect filled with Nova bone putty (GTa);
- (D) Socket defect filled with Bio Oss particulate bone graft (GTb)




Figure 1: Spider diagram depicting ISQ on buccal aspect.




Figure 2: Spider diagram depicting ISQ on mesial aspect.

Data Analysis




Figure 5: Bar chart comparing ISQ among different groups.




Figure 3: Spider diagram depicting ISQ on distal aspect.




Figure 4: Spider diagram depicting ISQ on lingual aspect.

Results

The non-parametric data was tested via Mann Whitney U test.

Initial torque value and ISQ						
GROUP		ITV	ISQ (Buccal)	ISQ (Mesial)	ISQ (Distal)	ISQ(Lingual)
Control (N=5)	Mean	35.00	54.00	55.80	55.60	51.80
	SD	.000	6.671	7.190	5.595	7.727
Defect and no graft (N=5)	Mean	20.00	38.60	36.20	39.00	37.80
	SD	.000	8.735	10.592	8.515	7.396
Defect and putty (N=5)	Mean	20.00	41.80	45.40	46.00	41.00
	SD	.000	5.020	9.864	10.440	5.339
Defect and particulate graft (N=5)	Mean	20.00	40.60	42.00	42.00	40.00
	SD	.000	4.615	3.808	4.848	9.618
Total	Mean	23.75	43.75	44.85	45.65	42.65
	SD	6.664	8.571	10.559	9.538	8.964

There was no statistically significant difference between the groups, however, highest ISQ values were observed in all putty groups.

Comparison among groups: Mann Whitney U test					
	GROUP	N	Mean Rank	Sum of Ranks	P value
ISQ (Buccal)	Defect and no graft	5	5.00	25.00	0.60
	Defect and putty graft	5	6.00	30.00	
	Defect and no graft	5	5.40	27.00	0.91
	Defect and particulate graft	5	5.60	28.00	
	Defect and putty graft	5	5.70	28.50	0.83
	Defect and particulate graft	5	5.30	26.50	
ISQ (Mesial)	Defect and no graft	5	4.10	20.50	0.14
	Defect and putty graft	5	6.90	34.50	
	Defect and no graft	5	4.90	24.50	0.52
	Defect and particulate graft	5	6.10	30.50	
	Defect and putty graft	5	6.40	32.00	0.34
	Defect and particulate graft	5	4.60	23.00	
ISQ (Distal)	Defect and no graft	5	4.40	22.00	0.25
	Defect and putty graft	5	6.60	33.00	
	Defect and no graft	5	4.90	24.50	0.52
	Defect and particulate graft	5	6.10	30.50	
	Defect and putty graft	5	6.40	32.00	0.34
	Defect and particulate graft	5	4.60	23.00	
ISQ (Lingual)	Defect and no graft	5	4.70	23.50	0.40
	Defect and putty graft	5	6.30	31.50	
	Defect and no graft	5	5.20	26.00	0.75
	Defect and particulate graft	5	5.80	29.00	
	Defect and putty graft	5	5.80	29.00	0.75
	Defect and particulate graft	5	5.20	26.00	

Conclusion

There was an increase in the stability of the implant with putty graft more than particulate graft or when no graft was placed in the defect. However, the outcome was not statistically significant. Further studies are needed to make a definitive conclusion.

Acknowledgement

We are sincerely thankful for the support of the Buhite-DiMino Corridor for Oral and Maxillofacial Health Foundation in the form of a grant which enabled me to attend this meeting.

References

1. Alkudmani H, AlJasser R, Andreana S. Is bone graft or guided bone regeneration needed when placing immediate dental implants? A systematic review. *Implant Dentistry*. Dec;26(6):936-944
2. Chen ST, Buser D (2009) Clinical and esthetic outcomes of implants placed in post extraction sites. *Int J Oral Maxillofac Implants* 24:186–217
3. Araújo MG, Wennström JL, Lindhe J (2006) Modeling of the buccal and lingual bone walls of fresh extraction sites following implant installation. *Clin Oral Implants Res* 17:606–614
4. Jones, A. A., Buser, D., Schenk, R., Wozney, J., & Cochran, D. L. (2006). The Effect of rhBMP-2 Around Endosseous Implants With and Without Membranes in the Canine Model. *Journal of Periodontology*, 77(7), 1184-1193. <https://doi.org/10.1902/jop.2006.050337>