

THE 41-YEAR MANAGEMENT OF A PATIENT WHO PRESENTED WITH EXTENSIVE BONE LOSS AT THE AGE OF 23



6/18/84: 23 Y.O.
INITIALLY
PRESENTED WITH
>50% BONE LOSS
Perio Treatment Plan:
scaling / root planing
/ occlusal adjustments
with 2- or 3-month
recalls



Picture used with patient permission

Thomas J. Kepic, DDS, MSD, Periodontist
Charles J. Goodacre, DDS, MSD, Prosthodontist

3/16/88, age 27




4 years after initial treatment



6/20/94, age 33

Example of annual perio chart

[illegible]






2017: 27 years later


Sequentially removed mandibular teeth
due to hypermobility, inability to chew
effectively, and discomfort

3/19/20 Follow Up Series

(Implants were placed 7/26/13)



Prosthesis designed with indentation for effective oral hygiene



What Factors Made This Treatment Successful For 39+ Years?

- Collaboration in planning and treatment
- Patient commitment to regular maintenance (2- or 3-month re-care)
- Prosthesis design for oral hygiene access

What would **YOUR** treatment plan be if this 23-year-old patient presented to your office today?

- Hirschfeld L, Wasserman B. A long-term survey of tooth loss in 600 treated periodontal patients. J Periodontol. 1978; 49: 225-237.
- McGuire MK, Nunn ME. Prognosis versus actual outcome. II. The effectiveness of clinical parameters in developing an accurate prognosis. J Periodontol. 1996; 67: 658-665.
- Rattu V, Raindi D, Antonoglou G, Nibali L. Prevalence of stable and successfully treated periodontitis subjects and incidence of subsequent tooth loss within supportive periodontal care: A systematic review with meta-analyses. J Clin Periodontol. 2023 Oct;50(10):1371-1389.
- 19 additional references available upon request