


Cervantes Manzano Z. V. DDS ¹ ; Nieto Ramírez J. A. DDS ²; Manisekaran R. DSc ²

¹ Postgraduate degree in periodontics and oral implantology student. Escuela Nacional de Estudios Superiores Unidad León UNAM


² Postgraduate degree in periodontics and oral implantology academic Escuela Nacional de Estudios Superiores Unidad León UNAM

Introduction


During dental extraction procedures, it is common to encounter intraoperative complications, with root fracture being a notable one. Retained root fragments that go undetected during implant placement can potentially serve as a source of pathogens in the etiology of periimplantitis. The purpose is to present a surgical protocol the simultaneous placement of a dental implant and the extraction of retained root.




Tissues analysis



Reteined root



Vestibular distance



Implant planing

Implant 4.6	
Implant type	Tapered plus conic
Conection type	Internal hexagonal
Body diameter	4.6 mm
Heigth	12 mm


Implant feature

Case presentation


A 49-year-old female patient presented for dental implant placement at the site of a missing tooth (4.6). Clinical analysis of the partially edentulous ridge revealed a thick phenotype and an adequate amount of keratinized tissue. Preoperative assessment of hard tissues was conducted through cone-beam computed tomography (CBCT), enabling analysis of dimensions and the presence of a retained root on the vestibular side of the ridge was identified.




Lateral window Technique



Implant placement



Healing abutment



Final radiography

Methods

A semi-restrictive analog guide was used, crestal and two vertical incisions over the edentulous ridge were made. Full-thickness flaps were used on the vestibular and lingual. The drilling sequence was initiated, and before implant placement, an osteotomy was performed through a facial lateral window to remove the retained root with an apical elevator. The dental implant was placed with an initial torque of 40 N/cm². Finally, a healing abutment was placed, and the flaps were approximated with simple interrupted sutures.

Results

Over a 12-week follow-up period, clinical evaluation revealed the absence of edema, erythema, suppuration, and bleeding. We observed adequate growth of keratinized tissue. Radiographically, in the non-critical bone defect (Fig. 2D) the filling of the same was observed (Fig. 3D).

Conclusion

This approach of extraction of a retained root, using the vestibular lateral window technique, provides an effective minimally invasive solution for the successful management of similar cases in order to decrease the probability of peri-implantitis.


References



Occlusal view of 12 weeks



Lateral view of 12 weeks



Direc pick-up coping