

A Treatment Strategy to Achieve the Complete Healing from Severe Peri-Implantitis

Eri Daikoku¹, Kyoko Takafuji¹, Kazuhiro Kon², Hisatomo Kondo¹

¹Aichi Gakuin University ²Iwate Medical University

Introduction

Peri-implantitis is a biological complication of the dental implant, and the absolute number of peri-implantitis is supposed to be increasing due to the recent spread of implant therapy. Currently, various studies have been conducted on the pathology and clinical diagnosis of peri-implantitis. However, the effective method has not yet been established, and this issue remains a major challenge in achieving long-term implant success. Here, we report that the clinically favorable result for peri-implantitis was obtained by the combination of mechanical cleaning with a titanium wire brush and autogenous bone grafting.

Diagnosis

Chief complaint of the patient: Discomfort of soft tissue around the implant.


History: The dental implant of molar region had been placed at another hospital more than five years ago, pus was found to be draining from the pocket, and X-ray photos revealed severe bone defect. The patient was diagnosed with peri-implantitis.




2013/08/28 First visit




2014/05/01 Preoperative X-ray photos




Pus was observed



Probing depth was 5 mm



Probing after removal of superstructure






Method and Results

2014/05/01



Decontamination with titanium wire brush



Autologous bone was harvested from the buccal side of molar region





Autogenous bone graft




6 months later

2014/11/13 2nd Surgery



Bone formation was confirmed

The postoperative soft tissue was kept in good shape, and the grafted bone around the implant was also restored. Although about ten years have passed since the final restoration was placed on, bone resorption was not observed in X-ray examinations at the regular checkup. Also, the patient is fully satisfied with the function and aesthetics.



2014/12/25 Provisional restoration



2015/02/24 Final restoration



2016/01/05 One year




2018/08/29 Four years




2023/10/04 Eight years

Conclusion

Complete healing from peri-implantitis was accomplished with recovery from peri-implant bone loss. This implant has been maintained in good condition for ten years, suggesting that this method is effective to recover peri-implantitis debridement with titanium wire brush and autogenous bone graft is one of the effective treatment option for peri-implantitis.



Autogenous bone grafting



Titanium wire brush

