

Non-abrasive peri-implantitis treatment to maximize titanium surface cytocompatibility

Rebecca Vaeth DDS 1; Ioanna Tsolaki DMD, MS, PhD 2; Georgios Kotsakis DDS, MS 3
 Advanced Graduate Dental Education, Division of Periodontics, Rutgers School of Dental Medicine, Newark, New Jersey
 Department of Oral Biology & Clinical Research Center, Rutgers School of Dental Medicine, Newark, New Jersey

Introduction

It has been suggested that instrumentation of the implant surface results in release of titanium particles which may induce an inflammatory response¹. Decontamination of the implant surface using chemotherapeutic agents during treatment of peri-implantitis defects may minimize the release of these titanium particles and preserve the implant’s surface properties². Additionally, the use of adjunctive biological materials such as enamel matrix derivative (EMD) has been positively associated with implant survival³.


Methods and Materials

I am presenting a case where a peri-implantitis defect was treated using a non-abrasive approach which included removal of the surrounding granulation tissue, burnishing of the exposed titanium surface with a NaOCl solution, placement of allograft combined with EMD, and a connective tissue graft.


Results

At a 6 month follow up, radiographic presentation of the peri-implant defect appears to have bone fill. Clinically, the overlying soft tissues appears thicker.


Pre-op



Surgical Procedure



Post-op (6 months)



Conclusions


A peri-implantitis treatment method in which the implant surface is burnished with NaOCl and the defect was grafted with allograft mixed with EMD can halt the progression of peri-implantitis while re-establishing healthy bone levels. Further research with extended follow-up would be necessary to determine reproducibility and long-term effectiveness of this technique.

References

- ¹ **Eger et al.** *Scientific Report*; 7:39612. (2017)
- ² **Kotsakis et al.** *Journal of Periodontology*, 87: 809-819. (2016)
- ³ **Ished et al.** *Journal of Clinical Periodontology*, 45: 744–753. (2018)

Non-abrasive peri-implantitis treatment to maximize titanium surface cytocompatibility

Rebecca Vaeth DDS 1; Ioanna Tsolaki DMD, MS, PhD 2; Georgios Kotsakis DDS, MS 3
Advanced Graduate Dental Education, Division of Periodontics, Rutgers School of Dental Medicine, Newark, New Jersey
Department of Oral Biology & Clinical Research Center, Rutgers School of Dental Medicine, Newark, New Jersey




Non-abrasive peri-implantitis treatment to maximize titanium surface cytocompatibility

Rebecca Vaeth DDS 1; Ioanna Tsolaki DMD, MS, PhD 2; Georgios Kotsakis DDS, MS 3
Advanced Graduate Dental Education, Division of Periodontics, Rutgers School of Dental Medicine, Newark, New Jersey
Department of Oral Biology & Clinical Research Center, Rutgers School of Dental Medicine, Newark, New Jersey





Introduction

It has been suggested that instrumentation of the implant surface results in release of titanium particles which may induce an inflammatory response¹. Decontamination of the implant surface using chemotherapeutic agents during treatment of peri-implantitis defects may minimize the release of these titanium particles and preserve the implant's surface properties². Additionally, the use of adjunctive biological materials such as enamel matrix derivative (EMD) has been positively associated with implant survival³.

Pre-op



Surgical Procedure



Methods and Materials

The peri-implantitis defect was treated using a non-abrasive approach:

- 0.5mm submarginal incision and intrasulcular incision was made around the implant to remove granulation tissue with minimal contact to the implant surface. FTF was elevated on the buccal.
- A solution of 1 part of NaOCl to 10 parts sterile water was applied via gentle burnishing with gauze for 10 seconds
- The implant was irrigated with ample saline
- A combination of allograft with EMD was placed in the defect
- A SCTG was placed buccal to the implant
- Primary closure was achieved with 5-0 vicryl sutures
- Results were assessed radiographically after 6 months of healing

Non-abrasive peri-implantitis treatment to maximize titanium surface cytocompatibility

Rebecca Vaeth DDS 1; Ioanna Tsolaki DMD, MS, PhD 2; Georgios Kotsakis DDS, MS 3
Advanced Graduate Dental Education, Division of Periodontics, Rutgers School of Dental Medicine, Newark, New Jersey
Department of Oral Biology & Clinical Research Center, Rutgers School of Dental Medicine, Newark, New Jersey

Results

At a 6 month follow up, radiographic presentation of the peri-implant defect appears to have bone fill. Clinically, the overlying soft tissues appears thicker.


Conclusions

A peri-implantitis treatment method in which the implant surface is burnished with NaOCl and the defect was grafted with allograft mixed with EMD can halt the progression of peri-implantitis while re-establishing healthy bone levels. Further research with extended follow-up would be necessary to determine reproducibility and long-term effectiveness of this technique.

References

- ¹ **Eger et al.** *Scientific Report*; 7:39612. (2017)
- ² **Kotsakis et al.** *Journal of Periodontology*, 87: 809-819. (2016)
- ³ **Isehmed et al.** *Journal of Clinical Periodontology*, 45: 744–753. (2018)

Pre-op



Post-op (6months)

