

Management of a Retained Epidural Catheter in a Pregnant Patient

Kevin W. Tang¹, Mingzhao Pei¹, Aamod George^{1,2}, Antoine Anderson^{1,2}, Ming Xiong^{1,2}

1. Rutgers New Jersey Medical School 2. Department of Anesthesia and Perioperative Care

BACKGROUND

Epidural Analgesia Overview:

- Commonly used to manage pain during labor and delivery, maintaining patient consciousness.
- Suitable for women requiring prolonged pain relief during labor; also used for long-duration Caesarean sections with combined spinal-epidural methods.

Administration and Patient-Controlled Epidural Analgesia (PCEA):

- Catheter placement in the epidural space enables continuous anesthetic delivery during prolonged labor.
- A study on PCEA with 0.0625% bupivacaine and fentanyl showed 92% satisfaction among 294 women, with fewer than 10% reporting complications.


Complications of Epidural Analgesia:

- Common side effects: hypotension, headache, back pain.
- Rare adverse effects: infection, nerve damage, and exceptionally rare retained catheter fragments.
- Lower concentrations of anesthetics (e.g., 0.0625% bupivacaine) and lipophilic opioids (fentanyl, sufentanil) have decreased adverse events.

Retained Epidural Catheter Fragment – Rare Complication:

- Symptoms range from minor pain to severe complications, though cases are often asymptomatic.
- Incidence is undocumented; case reports guide management, particularly for complex extractions where symptoms or fragment location warrant surgical intervention.

CASE PRESENTATION



DISCUSSION

Literature Insights on Retained Catheter Management:

- Most cases are asymptomatic and do not require surgery; conservative management often sufficient.
- Cases with persistent pain or neurological symptoms may need neurosurgical intervention to prevent nerve compression or fibrosis.

Imaging and Safety Considerations:

- Immediate imaging (e.g., MRI, CT, or x-ray) advised upon catheter fracture detection; MRI may pose thermal risks with metal-reinforced tips.
- CT imaging used in this case due to metallic spring in catheter tip, prioritizing safety.

Prevention Techniques:

- Suggested techniques to avoid retention include gentle retraction, placing patients in lateral decubitus position, minimizing catheter insertion depth, and using low tension during removal.

Case Insights and Recommendations:

- Fracture likely due to multiple insertion attempts and increased traction.
- Importance of early imaging, understanding catheter type, and patient reassurance emphasized.
- Call for future research on the causes, incidence, and management protocols for retained catheters, as well as MRI compatibility for metal-reinforced tips.

REFERENCES

- [1] Hawkins, J.L. (2010) Epidural Analgesia for Labor and Delivery. New England Journal of Medicine, 362, 1503-1510. <https://doi.org/10.1056/nejmct0909254>
- [2] Author (2016) Practice Guidelines for Obstetric Anesthesia: An Updated Report by the American Society of Anesthesiologists Task Force on Obstetric Anesthesia and the Society for Obstetric Anesthesia and Perinatology. Anesthesiology, 124, 270-300.
- [3] Clivatti, J., Siddiqui, N., Goel, A., Shaw, M., Crisan, I. and Carvalho, J.C.A. (2013) Quality of Labour Neuraxial Analgesia and Maternal Satisfaction at a Tertiary Care Teaching Hospital: A Prospective Observational Study. Canadian Journal of Anesthesia, 60, 787-795. <https://doi.org/10.1007/s12630-013-9976-9>
- [4] Parnass, S.M. and Schmidt, K.J. (1990) Adverse Effects of Spinal and Epidural Anesthesia. Drug Safety, 5, 179-194. <https://doi.org/10.2165/00002018-199005030-00003>
- [5] Sharma, R., Setlur, R., Bhargava, A. and Vardhan, S. (2007) Walking Epidural: An Effective Method of Labour Pain Relief. Medical Journal Armed Forces India, 63, 44-46. [https://doi.org/10.1016/s0377-1237\(07\)80107-9](https://doi.org/10.1016/s0377-1237(07)80107-9)
- [6] Merrer, J., Bonnet, M., Blondel, B., Tafflet, M., Khoshnood, B., Le Ray, C., et al. (2021) Predictors of Incomplete Maternal Satisfaction with Neuraxial Labor Analgesia: A Nationwide Study. Anaesthesia Critical Care & Pain Medicine, 40, Article 100939. <https://doi.org/10.1016/j.accpm.2021.100939>