

AN UNUSUAL CASE OF RUPTURED INTRAMURAL ECTOPIC IN THE SECOND TRIMESTER

Dr Mai Khanh Kieu Xuan, Dr Narjes Fhelelboom, Dr Nandini Ravikumar, Dr Waleed Khattab, Prof. Sam Thomas, Prof Michael Gannon, Dr Majda Almshwt
Department of Obstetrics & Gynaecology, Mullingar Regional Hospital, Mullingar, Ireland



Feidhmeannacht na Seirbhíse Sláinte
Health Service Executive



Royal College of
Obstetricians &
Gynaecologists



INTRODUCTION

We report an unusual case where the diagnosis of a **ruptured intramural ectopic** at 15th gestation week was made intraoperatively.

DEFINITION	<p>Ectopic pregnancy remains one of the leading causes of maternal mortality worldwide. Ectopic pregnancy is considered unsustainable and potential life threatening in case of rupture.</p> <p>Intramural ectopic is described when the gestation sac is implanted within the myometrium separate from the uterus cavity, endometrium and the fallopian tubes.</p>
EPIDEMIOLOGY AETIOLOGY	<p>Intramural ectopic is one of the rarest forms that happens less than 1% of ectopic pregnancies.</p> <p>Aetiology of intramural ectopic remains debatable. Several risk factors such as uterine surgery including dilation and curettage, previous cesarean section, or myomectomy may have a contributing role.</p>
SYMPTOMS	<p>In general, the classic triad of a suspected ectopic pregnancy include: abdominal and/or pelvic pain, vaginal bleeding, and a positive pregnancy test. Syncope or hypotension will raise further alert for possible ruptured ectopic.</p>
DIAGNOSIS	<p>Diagnosis of ectopic pregnancy is confirmed with serial ultrasound scans and serial serum hCG in the first trimester.</p> <p>From the literature, for intramural ectopic the diagnosis is usually a combination of transvaginal ultrasound, MRI, hysteroscopy and histopathology.</p>
TREATMENT	<p>For non-rupture ectopic pregnancy, expectant management, medical management with methotrexate, and surgical management are the current options.</p> <p>Ruptured ectopic is a medical emergency therefore surgery is always needed. Preserved fertility is varied with different cases.</p>




CASE REPORT

A 42 year old female G5P1 (previous Caesarean section) presented to emergency department with sudden onset abdominal pain at 15 weeks pregnancy.



HISTORY

- She was known to have an **anterior uterine fibroid**, awaiting surgery but postponed because of this unplanned pregnancy.
- Her first ultrasound scan at 8 weeks was **inconclusive** due to the large fibroid and increased BMI.
- The subsequent booking scan at 12 weeks reported intrauterine pregnancy.



Gestational sac and fetus heartbeat detected at 10 gestation's week.



Anterior wall fibroid size: 4x6cm



EMERGENCY PRESENTATION

- Patient was in severe distress with tenderness all over her abdomen on slight movements and palpation. However she reports **no** vaginal bleeding or any pregnancy related symptoms.

- An informal bed-side ultrasound at the time revealed the known fibroid which obstruct the view significantly. Nevertheless, a single foetus was seen with active heart activity and a small amount of free fluid.
- Biochemistry blood results are reported all within normal range with no ongoing infection. Her Hb was stable at 13g/dL despite her pale looking appearance.




HOSPITALISATION EVENTS

- Despite escalation of analgesia, patient still suffered from persisted and worsened abdominal pain.
- A second set of blood test were repeated approximately 3 hours after admission; this time her Hb was noted to drop down to 8.5 g/dL while other biochemistry test remains unchanging.
- Consultation was made between specialties including surgical team, obstetrician team, medical team and anaesthetic team as patient started deteriorating with low BP of 70/40mmHg.
- A decision was made for emergency laparoscopy to identify cause of a suspected **internal bleeding**.



OPERATION FINDINGS

- As soon as the scope entered the abdominal cavity, hemoperitoneum was noted in abundance.
- The emergency laparoscopy procedure was converted to laparotomy the moment we identified foetus feet in the abdomen cavity with the first impression of uterine rupture.
- However, the placenta and remnant of sac was removed without entering the uterine cavity. Intramural ectopic rupture was then diagnosed during operation.



Schematic drawing of sagittal view of uterus depicting the location of intramural ectopic sac in relation to the anterior wall fibroid.

(Drawn by Dr Mai Khanh Kieu Xuan)



DISCUSSION

The patient's moderate-sized fibroid have misguided the intramural pregnancy to look like an intrauterine one at the early stage. In addition, her lack of classical ectopic symptoms and the initial normal blood result can be deceiving. Therefore, our take home message which we learned from this case is:

- Despite the normal initial lab result, it is always sensible to reassess when patient's symptoms persist or worsen. Repeated investigations may be warranted for a proper monitoring from the re-assessment.

REFERENCES: 1. Molinaro TA, Barnhart KT. Ectopic pregnancies in unusual locations. Semin Reprod Med 2007;25:123–30.

2. Chen, Xiaoran et al. "Intramural Ectopic Pregnancy: Clinical Characteristics, Risk Factors for Uterine Rupture and Hysterectomy." Frontiers in medicine vol. 8 769627. 28 Oct. 2021, doi:10.3389/fmed.2021.769627