

EP.0176. Encountering placenta accreta in first trimester missed miscarriage- A case report

OBJECTIVE: To emphasize the rare encounter of morbidly adherent placenta in first trimester surgical evacuation for missed miscarriage that can risk life of patient.

CASE REPORT: A 33 years old G2P1+0 with previous history of LSCS one year back due to symptomatic placenta previa at 36 wks presented in O&G clinic with ultrasound diagnosis of missed miscarriage at 8 wks.

- She underwent medical T.O.P but as had no response to misoprostol, she was booked for suction evacuation under G.A.
- With the introduction of plastic suction catheter tube, the visible deterioration of pulse and blood pressure was noticed.

An intra operative scan revealed massive intra peritoneal hemorrhage.

- An emergency laparotomy and code blue instituted with initiation of massive obstetric hemorrhage protocol from O.T.
- Per operative 2 to 3 L of blood was removed, uterus and adnexal areas snugly packed with surgical packs and aortic compression with the fist of operating surgeon done.
- When patient's vitals stabilized with blood transfusion and i.v fluid resuscitation and she recovered from hypovolemic hemorrhagic shock, the abdominal packs removed.
- A bluish discolored area of 1.5x2cm cm over uterine isthmus at the site of previous scar was noticed. The uterine walls were intact. A small incision was given over the bluish area on the anterior uterine wall and placental tissue scooped out digitally. From same incision uterine cavity was sucked out for any RPOCs.
- Uterine incision was closed in 2 layers with vicryl no 2/0.

