

Labor outcomes in comprehensive COVID 19 tested population in tertiary care facility

Dr. Ushna Ahmed, Dr. Farheen Yousuf, Dr. Irfan Sheikh, Miss. Ghulam Kubra, Department of Obstetrics and Gynecology, Aga Khan University Hospital, Karachi, Pakistan

Abstract	Results	Discussion																																																																																																																																																									
Early detection of SARS CoV 2 status in pregnant patients is crucial, this allows for risk appropriate, judicious use of personal protective equipment (PPE), with timely and appropriate neonatal testing and rooming practices. It also provides safe environment for the neonate, family members, other patients, and health staff.	Maternal Characteristics In total 2887 women who delivered,1918 women were subjected to COVID-19 testing. 5.52% tested covid-positive and 94.4% tested negative. 7 were symptomatic and 93.4% were asymptomatic, most commonly reported sore throat. A significant difference of BMI was found in both the groups (p-value 0.001), women with Covid-19 positive had higher BMI. More women with covid positive were admitted for labor indication and for elective lower segment cesarean section. GDM was found to be higher among women with covid positive than negative. No association was found in other risk factor amongst those who tested positive and negative. <table><tr><td>Pregestational Hypertension</td><td>1 (0.9%)</td><td>83 (4.6%)</td><td>0.085</td></tr><tr><td>Asthma</td><td>1 (0.9%)</td><td>21 (1.2%)</td><td>0.999</td></tr><tr><td>Smoking</td><td>0 (0%)</td><td>2 (0.1%)</td><td>0.999</td></tr></table> Data are presented as median [25-75 percentile], n (%). Mann-Whitney U for median, and Chi-square with a Fisher Exact test (<=5) for n (%).* Denotes significance Obstetric outcome Women with covid positive had more c-section than covid-negative. In indications, non-reassuring fetal status was found to be higher among covid positive women than negative. Scheduled repeat cesarean sections were more among Covid-19 positive women than negative. Cesarean sections for mal-presentation were more in the Covid-19 positive group than negative. Non-progress of labor was less in Covid-19 positive women. Neonatal outcomes <table><tr><td>Neonatal outcomes</td><td>SARS-CoV-2 RT-PCR Positive (n=106)</td><td>SARS-CoV-2 RT-PCR Negative (n=1812)</td><td>P-value</td></tr><tr><td>Birth weight</td><td>2.9[2.5-3.2]</td><td>2.9[2.6-3.1]</td><td>0.996</td></tr><tr><td>Live birth</td><td>104 (98.1%)</td><td>1786 (98.6%)</td><td>0.665</td></tr><tr><td>Still birth</td><td>1 (0.9%)</td><td>1 (0.9%)</td><td>0.108</td></tr><tr><td>TUD</td><td>1 (0.9%)</td><td>25 (1.4%)</td><td>0.999</td></tr><tr><td>Preterm birth</td><td>30 (28.3%)</td><td>368 (20.3%)</td><td>0.049*</td></tr><tr><td>APGAR Score at 1 min (Missing=28)</td><td></td><td></td><td></td></tr><tr><td><7</td><td>0 (0%)</td><td>20 (1.1%)</td><td>0.623</td></tr><tr><td>≥7</td><td>104 (98.1%)</td><td>1766 (97.5%)</td><td>0.999</td></tr><tr><td>APGAR Score at 5 min (Missing=28)</td><td>104 (98.1%)</td><td>1786 (98.6%)</td><td>0.665</td></tr></table> Data are presented as median [25-75 percentile], n (%). Mann-Whitney U for median, and Chi-square with a Fisher Exact test (<=5) for n (%).* Denotes significance. Table I (Maternal characteristics stratified by SARS-CoV-2 RT-PCR result) <table><tr><th>Maternal Characteristics</th><th>SARS-CoV-2 RT-PCR Positive (n=106)</th><th>SARS-CoV-2 RT-PCR Negative (n=1812)</th><th>P-value</th></tr><tr><td>Age</td><td>29.5[26-35]</td><td>28[27-32]</td><td>0.083</td></tr><tr><td>BMI</td><td>29.5[27-32]</td><td>28[26-30]</td><td>0.001*</td></tr><tr><td>Multiple gestation</td><td>0 (0%)</td><td>8 (0.4%)</td><td>0.999</td></tr><tr><td>Gestational age at admission</td><td>38 [36-39]</td><td>38[37-39]</td><td>0.522</td></tr><tr><td>Gravida</td><td></td><td></td><td></td></tr><tr><td>Primi</td><td>35(33%)</td><td>669(36.9%)</td><td rowspan="3">0.720</td></tr><tr><td>Multi</td><td>60(56.6%)</td><td>967(53.4%)</td></tr><tr><td>Grand Multi</td><td>11(10.4%)</td><td>176(9.7%)</td></tr><tr><td>Booking Status</td><td></td><td></td><td></td></tr><tr><td>Booked</td><td>96(90.6%)</td><td>1648(90.9%)</td><td rowspan="2">0.89</td></tr><tr><td>Unbooked</td><td>10(9.4%)</td><td>164(9.1%)</td></tr><tr><td>Indication for admission</td><td></td><td></td><td></td></tr><tr><td>Labour indication</td><td>44(41.5%)</td><td>609(33.6%)</td><td>0.095</td></tr><tr><td>Fetal indication</td><td>13(12.3%)</td><td>382(21.1%)</td><td>0.026*</td></tr><tr><td>Elective LSCS</td><td>26 (24.5%)</td><td>314 (17.3%)</td><td>0.067</td></tr><tr><td>Elective IOL</td><td>17 (16%)</td><td>441 (24.3%)</td><td>0.060</td></tr><tr><td>PPROM</td><td>2 (1.9%)</td><td>64 (3.5%)</td><td>0.581</td></tr><tr><td>COVID like symptoms</td><td>0 (0%)</td><td>2 (0.1%)</td><td>0.999</td></tr><tr><td>Others</td><td>4 (3.8%)</td><td>0 (0%)</td><td>0.000*</td></tr><tr><td>Comorbidities</td><td></td><td></td><td></td></tr><tr><td>GDM</td><td>33 (31.1%)</td><td>371 (20.5%)</td><td>0.009*</td></tr><tr><td>PIH/ Preeclampsia</td><td>11 (10.4%)</td><td>238 (13.1%)</td><td>0.412</td></tr><tr><td>Hyper/hypothyroidism</td><td>5 (4.7%)</td><td>102 (5.6%)</td><td>0.691</td></tr><tr><td>Risk Factors</td><td></td><td></td><td></td></tr><tr><td>Pregestational diabetes</td><td>3 (2.8%)</td><td>52 (2.9%)</td><td>0.999</td></tr></table>	Pregestational Hypertension	1 (0.9%)	83 (4.6%)	0.085	Asthma	1 (0.9%)	21 (1.2%)	0.999	Smoking	0 (0%)	2 (0.1%)	0.999	Neonatal outcomes	SARS-CoV-2 RT-PCR Positive (n=106)	SARS-CoV-2 RT-PCR Negative (n=1812)	P-value	Birth weight	2.9[2.5-3.2]	2.9[2.6-3.1]	0.996	Live birth	104 (98.1%)	1786 (98.6%)	0.665	Still birth	1 (0.9%)	1 (0.9%)	0.108	TUD	1 (0.9%)	25 (1.4%)	0.999	Preterm birth	30 (28.3%)	368 (20.3%)	0.049*	APGAR Score at 1 min (Missing=28)				<7	0 (0%)	20 (1.1%)	0.623	≥7	104 (98.1%)	1766 (97.5%)	0.999	APGAR Score at 5 min (Missing=28)	104 (98.1%)	1786 (98.6%)	0.665	Maternal Characteristics	SARS-CoV-2 RT-PCR Positive (n=106)	SARS-CoV-2 RT-PCR Negative (n=1812)	P-value	Age	29.5[26-35]	28[27-32]	0.083	BMI	29.5[27-32]	28[26-30]	0.001*	Multiple gestation	0 (0%)	8 (0.4%)	0.999	Gestational age at admission	38 [36-39]	38[37-39]	0.522	Gravida				Primi	35(33%)	669(36.9%)	0.720	Multi	60(56.6%)	967(53.4%)	Grand Multi	11(10.4%)	176(9.7%)	Booking Status				Booked	96(90.6%)	1648(90.9%)	0.89	Unbooked	10(9.4%)	164(9.1%)	Indication for admission				Labour indication	44(41.5%)	609(33.6%)	0.095	Fetal indication	13(12.3%)	382(21.1%)	0.026*	Elective LSCS	26 (24.5%)	314 (17.3%)	0.067	Elective IOL	17 (16%)	441 (24.3%)	0.060	PPROM	2 (1.9%)	64 (3.5%)	0.581	COVID like symptoms	0 (0%)	2 (0.1%)	0.999	Others	4 (3.8%)	0 (0%)	0.000*	Comorbidities				GDM	33 (31.1%)	371 (20.5%)	0.009*	PIH/ Preeclampsia	11 (10.4%)	238 (13.1%)	0.412	Hyper/hypothyroidism	5 (4.7%)	102 (5.6%)	0.691	Risk Factors				Pregestational diabetes	3 (2.8%)	52 (2.9%)	0.999	The prevalence of COVID-19 was found in 5.5 % in our data set while other studies reported the detection rate of between 0.5-10.5% on universal screening. Majority of them were asymptomatic while only 6.6 % had clinical symptoms suggestive of COVID 19. The ample literature available to verifies it, and emphasizes the need for generalized screening of all pregnant women to identify and isolate them to avoid spread of the disease. The common symptoms in our study were sore throat and cough. Other studies had reported fever as the commonest clinical presentation 62.9% followed by cough in 36.8% and sore throat in 22.6%. COVID-19 positive women had statistically significant propensity towards high BMI, which was also seen in a study, reporting 60.5 %of positive women were obese . Gestational diabetes mellitus was significantly higher in COVID-19 positive women, followed by hypertension (31.1% and 10.4 % respectively), which was reflected by a study from Stockholm. COVID-19 positive women were found to be more prone to developing hypertensive disorders in pregnancy, with the incidence being 34.2%. Other studies conducted under similar lines do not support preference towards a single major reason for admission, as suggested in our study. One of the pioneer studies suggested an increased cesarean delivery rate among women. A study from Pakistan also exhibit 78.5% rate of C-section in with COVID-19. A study from New York, suggested no increase in the rate of cesarean section amongst positive women. The neonatal outcome has generally been similar and satisfactory for both positive and negative tested women. This has been speculated by several observational studies and meta-analysis, which suggest no differences between the neonatal outcomes of COVID-19 positive and negative women. Vertical transmission was not reported in majority of studies, however a meta-analysis by Matar’R et.al found two cases of fetal Covid positive. This is the first of its kind study to be conducted in a tertiary care hospital form Karachi, Pakistan, to generate local data regarding the prevalence of COVID-19 in obstetric population, as well as the maternal and fetal characteristics of COVID-19 positive and negative parturient in Pakistan. The limitation of this study was conducted in a single tertiary care private hospital of Karachi; therefore, the results may vary and not be applicable to the rest of the public center hospitals of this country.
Pregestational Hypertension	1 (0.9%)	83 (4.6%)	0.085																																																																																																																																																								
Asthma	1 (0.9%)	21 (1.2%)	0.999																																																																																																																																																								
Smoking	0 (0%)	2 (0.1%)	0.999																																																																																																																																																								
Neonatal outcomes	SARS-CoV-2 RT-PCR Positive (n=106)	SARS-CoV-2 RT-PCR Negative (n=1812)	P-value																																																																																																																																																								
Birth weight	2.9[2.5-3.2]	2.9[2.6-3.1]	0.996																																																																																																																																																								
Live birth	104 (98.1%)	1786 (98.6%)	0.665																																																																																																																																																								
Still birth	1 (0.9%)	1 (0.9%)	0.108																																																																																																																																																								
TUD	1 (0.9%)	25 (1.4%)	0.999																																																																																																																																																								
Preterm birth	30 (28.3%)	368 (20.3%)	0.049*																																																																																																																																																								
APGAR Score at 1 min (Missing=28)																																																																																																																																																											
<7	0 (0%)	20 (1.1%)	0.623																																																																																																																																																								
≥7	104 (98.1%)	1766 (97.5%)	0.999																																																																																																																																																								
APGAR Score at 5 min (Missing=28)	104 (98.1%)	1786 (98.6%)	0.665																																																																																																																																																								
Maternal Characteristics	SARS-CoV-2 RT-PCR Positive (n=106)	SARS-CoV-2 RT-PCR Negative (n=1812)	P-value																																																																																																																																																								
Age	29.5[26-35]	28[27-32]	0.083																																																																																																																																																								
BMI	29.5[27-32]	28[26-30]	0.001*																																																																																																																																																								
Multiple gestation	0 (0%)	8 (0.4%)	0.999																																																																																																																																																								
Gestational age at admission	38 [36-39]	38[37-39]	0.522																																																																																																																																																								
Gravida																																																																																																																																																											
Primi	35(33%)	669(36.9%)	0.720																																																																																																																																																								
Multi	60(56.6%)	967(53.4%)																																																																																																																																																									
Grand Multi	11(10.4%)	176(9.7%)																																																																																																																																																									
Booking Status																																																																																																																																																											
Booked	96(90.6%)	1648(90.9%)	0.89																																																																																																																																																								
Unbooked	10(9.4%)	164(9.1%)																																																																																																																																																									
Indication for admission																																																																																																																																																											
Labour indication	44(41.5%)	609(33.6%)	0.095																																																																																																																																																								
Fetal indication	13(12.3%)	382(21.1%)	0.026*																																																																																																																																																								
Elective LSCS	26 (24.5%)	314 (17.3%)	0.067																																																																																																																																																								
Elective IOL	17 (16%)	441 (24.3%)	0.060																																																																																																																																																								
PPROM	2 (1.9%)	64 (3.5%)	0.581																																																																																																																																																								
COVID like symptoms	0 (0%)	2 (0.1%)	0.999																																																																																																																																																								
Others	4 (3.8%)	0 (0%)	0.000*																																																																																																																																																								
Comorbidities																																																																																																																																																											
GDM	33 (31.1%)	371 (20.5%)	0.009*																																																																																																																																																								
PIH/ Preeclampsia	11 (10.4%)	238 (13.1%)	0.412																																																																																																																																																								
Hyper/hypothyroidism	5 (4.7%)	102 (5.6%)	0.691																																																																																																																																																								
Risk Factors																																																																																																																																																											
Pregestational diabetes	3 (2.8%)	52 (2.9%)	0.999																																																																																																																																																								
Introduction Pakistan reported its first two confirmed cases of COVID-19 on February 26th, 2020 . Maternity health care units continue to provide care to its regular volume of patients. Generalized testing of all patients presenting to obstetrical units was proposed, and proved necessary. Following the initiation of universal screening programs by Sutton et.al, data on COVID-19 in pregnancy started emerging Recent estimates suggest <1% of asymptomatic patients have tested positive since initiating universal screening. Some studies suggested that pregnant women with COVID-19-related symptoms have a high rate of severe disease and preterm birth. It is prudent for obstetricians to be particularly vigilant when caring for such patients who become symptomatic. The Aga Khan University Hospital started testing women booked for elective procedures in obstetrics and gynecology for SARS CoV 2 in May 2020 Using data from these records our study aims to determine the prevalence of SARS CoV 2 amongst obstetrical patients presenting to obstetrical units, being subjected to generalized screening Also determine the characteristics of the disease amongst these women, its evolution during the hospital stay, differences in pregnancy outcomes between affected and unaffected females	Discussion																																																																																																																																																										
Objectives To determine the prevalence of SARS CoV 2 amongst obstetrical patients presenting to obstetrical units, being subjected to generalized screening. The study also aims to determine the characteristics of the disease amongst these women, its evolution during the hospital stay, differences in pregnancy outcomes between affected and unaffected females.	Conclusion There was low prevalence of COVID-19 among pregnant women, and mainstream were asymptomatic, these women tend to be carriers, and can spread the virus It is wise to implement such generalized COVID testing in the obstetric population																																																																																																																																																										
Materials and Methods This prospective observational study was conducted at the Department of Obstetrics and Gynecology, Aga Khan University hospital, Karachi, between the months of June and November 2020. All women >24 weeks of pregnancy presented to labour room pertaining to admission or delivery, underwent generalized screening for COVID 19 PCR along with symptoms related to it were and maternal characteristics were recorded. These women were observed during course of hospital stay. On bases of Covid-19 status they had been divided into Covid-19 positive and negative groups. Data collected on maternal characteristics that include maternal age, BMI, parity, gestational age at presentation, booking status, indication for admission, comorbidities, and antenatal risk factors. Data was collected on obstetrical outcome, mode of delivery, and indication for a cesarean section. For neonates, data regarding their birth weight, live birth, still birth or an intrauterine fetal demise, pre-term birth, and APGAR score at 1 and 5 minutes were collected	References: - Abid K, Bari YA, Younas M, Tahir Javid S, Imran A. Progress of COVID-19 Epidemic in Pakistan. Asia Pac J Public Health. 2020;32(4):154-156 doi:10.1177/1010539520927259. - Herraiz I, Folgueira D, Villalain C, Forcén L, Delgado R, Galindo A. Universal screening for SARS-CoV-2 before labor admission during Covid-19 pandemic in Madrid. J Perinat Med. 2020;48(9):981-984 doi: 10.1515/jpm-2020-0236. - Sutton D, Fuchs K, D’alton M, Goffman D. Universal screening for SARS-CoV-2 in women admitted for delivery. The New England journal of medicine. 2020;382(22):2163-2164 doi:10.1056/NEJMc2009316. - Chen L, Li Q, Zheng D, Jiang H, Wei Y, Zou L, et al. Clinical Characteristics of Pregnant Women with Covid-19 in Wuhan, China. The New England journal of medicine. 2020;382(25):e100 doi:10.1056/NEJMc2009226. - Prabhu M, Cagino K, Matthews KC, Friedlander RL, Glynn SM, Kubiak JM, et al. Pregnancy and postpartum outcomes in a universally tested population for SARS-CoV-2 in New York City: a prospective cohort study. BJOG. 2020;127(12):1548-1556 doi:10.1111/1471-0528.16403. - Fassett MJ, Lurvey LD, Yasumura L, Nguyen M, Colli JJ, Volodarskiy M, et al. Universal SARS-Cov-2 Screening in Women Admitted for Delivery in a Large Managed Care Organization. Am J Perinatol. 2020;37(11):1110-1114 doi:10.1055/s-0040-1714060																																																																																																																																																										

The prevalence of COVID-19 was found in 5.5 % in our data set while other studies reported the detection rate of between 0.5-10.5% on universal screening. Majority of them were asymptomatic while only 6.6 % had clinical symptoms suggestive of COVID 19. The ample literature available to verifies it, and emphasizes the need for generalized screening of all pregnant women to identify and isolate them to avoid spread of the disease.

The common symptoms in our study were sore throat and cough. Other studies had reported fever as the commonest clinical presentation 62.9% followed by cough in 36.8% and sore throat in 22.6%. COVID-19 positive women had statistically significant propensity towards high BMI, which was also seen in a study, reporting 60.5 %of positive women were obese . Gestational diabetes mellitus was significantly higher in COVID-19 positive women, followed by hypertension (31.1% and 10.4 % respectively), which was reflected by a study from Stockholm. COVID-19 positive women were found to be more prone to developing hypertensive disorders in pregnancy, with the incidence being 34.2%. Other studies conducted under similar lines do not support preference towards a single major reason for admission, as suggested in our study.

One of the pioneer studies suggested an increased cesarean delivery rate among women. A study from Pakistan also exhibit 78.5% rate of C-section in with COVID-19. A study from New York, suggested no increase in the rate of cesarean section amongst positive women.

The neonatal outcome has generally been similar and satisfactory for both positive and negative tested women. This has been speculated by several observational studies and meta-analysis, which suggest no differences between the neonatal outcomes of COVID-19 positive and negative women. Vertical transmission was not reported in majority of studies, however a meta-analysis by Matar’R et.al found two cases of fetal Covid positive.

This is the first of its kind study to be conducted in a tertiary care hospital form Karachi, Pakistan, to generate local data regarding the prevalence of COVID-19 in obstetric population, as well as the maternal and fetal characteristics of COVID-19 positive and negative parturient in Pakistan. The limitation of this study was conducted in a single tertiary care private hospital of Karachi; therefore, the results may vary and not be applicable to the rest of the public center hospitals of this country.