

Utilizing mHealth for implementing the WHO ANC model: A pilot study

S Soni¹, KA Sharma¹, V Perumal¹, K Yadav², S Singhal¹

¹Department of Obstetrics and Gynaecology, All India Institute of Medical Sciences, New Delhi-110029

²Centre for Community Medicine, All India Institute of Medical Sciences, New Delhi-110029



Introduction

- ❖ Increasing focus on the content and quality of care
- ❖ New antenatal care model with 8 “contacts” and objectives to ensure a “positive pregnancy experience”
- ❖ Respectful, individualized and person-centred care at each visit
- ❖ Interventions in the model addressed under 5 domains

WHO FANC Model	2016 WHO ANC Model
First Trimester	
Visit 1 : 8-12 weeks	Contact 1 : up to 12 weeks
Second Trimester	
Visit 2 : 24-26 weeks	Contact 2: 20 weeks Contact 3 : 26 weeks
Third Trimester	
Visit 3 : 32 weeks	Contact 4 : 30 weeks Contact 5 : 34 weeks
Visit 4 : 36-38 weeks	Contact 6 : 36 weeks Contact 7 : 38 weeks Contact 8 : 40 weeks

Patient Registration

Upto 14th weeks

Date: APR 19, 2022

POG: 1 WEEKS

Patient Name: Tripti Singh

PREGNANCY STAGE

History: ☐ Fever, ☐ Rash

Chest/CVS: Normal, Not Done, Abnormal

P/A: Nil

Others: Details

Counselling regarding Birth Plan: Yes, No

Birth Companion:

Place of delivery:

Mode of delivery:

Past Visits

High Risk

Antenatal Clinic Schedule activity recaller

- History (visit specific)
- Examination (visit specific)
- Investigations (visit specific):
- Advice (medications /diet/exercise/specific treatment/danger signs)
- Planning delivery (Mode, time, and preparations required)
- Advice on minor ailments encountered in pregnancy

Number of Visits


Number of Visits	Cases (n=85) (%)	Controls (n=73) (%)	p value
0-4	16 (18.8)	45 (61.6)	p<0.001
5-8	65 (76.5)	24 (32.9)	
>8	4 (4.7)	4 (5.5)	

Quality of Care

Quality Scores	Cases	Controls	p value
History	8.26 ± 2.26	3.57 ± 2.30	<0.001
Examination	4.26 ± 1.03	3.66 ± 1.09	0.001
Counselling	6.09 ± 1.65	4.33 ± 1.69	<0.001

Antenatal Investigations

Antenatal Investigations	Cases (n=85) (%)	Controls (n=73) (%)	p value
Hemogram	85 (100)	68 (93.2)	0.020
Urine	72 (92.9)	65 (89)	0.415
USG	76 (89.4)	41 (56.2)	0.0001




Aims and Objectives

- ❖ To compare the quality of antenatal care delivered by the use of a web-based mobile application compared to the standard care protocol as assessed by the content of care

Methodology


- ❖ Study Design : Multicentric prospective open-label RCT
- ❖ Study Site: Sub-district hospital and primary health centres in North India
- ❖ Study Duration: December 2019 to December 2021
- ❖ Randomization: Computer generated random allocation tables

Development of the app



All women booked before 20 weeks, possessing smartphone or having access to one and willing to give consent were included

Results



Baseline demographic variables were comparable

Domains	Components of care	Cases (n=85) (%)	Controls (n=73) (%)	p value
History Obtained	Medical History	80 (94.1)	22 (30.1)	0.0001
	Surgical History	78 (91.8)	10 (13.7)	0.0001
	Drugs	78 (91.8)	64 (87.7)	0.437
	Tobacco Use	77 (90.6)	21 (28.8)	0.0001
	Smoking	77 (90.6)	23 (31.5)	0.0001
	Alcohol	76 (89.4)	22 (30.1)	0.0001
	IPV	76 (89.4)	14 (19.1)	0.0001
Physical Examination	Fetal movement	83 (97.6)	66 (90.4)	0.082
	Height	85 (100)	70 (95.9)	0.096
	Weight	85 (100)	72 (98.6)	0.462
	Blood Pressure	80 (94.1)	66 (90.4)	0.549
	Fundal height	56 (65.9)	26 (35.6)	0.0001
	FHS	56 (65.9)	26 (35.6)	0.0001
	IFA and calcium supplementation	82 (96.5)	69 (94.5)	0.705
Patient Counselling	Dietary	80 (94.1)	38 (52)	0.0001
	Exercise	79 (92.9)	34 (46.6)	0.0001
	Tetanus	83 (97.6)	65 (89)	0.045
	Breastfeeding	44 (51.7)	12 (16.4)	0.0001
	Pain relief in labour	19 (22.3)	8 (11)	0.058

Using the PreMAPEQ questionnaire, higher mean satisfaction score was seen among the cases

Discussion

- ❖ Utility of mHealth system and impact on quality of antenatal care demonstrated by other studies also (Benski et al, McNabb et al)
- ❖ Comprehensive inclusion of a larger number of parameters and emphasis on indicators affecting quality
- ❖ However, larger sample size required for impact on perinatal outcomes

Conclusion

- ❖ mHealth technology has an immense potential to improve the quality of healthcare
- ❖ Significant improvement seen in this study with the use of mobile technology in parameters such as communication with the patients and counselling provided, resulting in a “positive pregnancy experience”

References

- World Health Organization. WHO Recommendations on Antenatal Care for a Positive Pregnancy Experience. WHO Guidelines Approved by the Guidelines Review Committee. WHO Recommendations on Antenatal Care for a Positive Pregnancy Experience. World Health Organization; 2016.
- Villar J, Ba'aqeel H, Piaggio G, Lumbiganon P, Miguel Belizán J, Farnot U, et al. WHO antenatal care randomised trial for the evaluation of a new model of routine antenatal care. *Lancet*. 2001 May 19;357(9268):1551–64.
- Benski AC, Schmidt NC, Viviano M, Stancanelli G, Soaroby A, Reich MR. Improving the Quality of Antenatal Care Using Mobile Health in Madagascar: Five-Year Cross-Sectional Study. *JMIR mHealth uHealth*. 2020 Jul 8;8(7):e18543.
- McNabb M, Chukwu E, Ojo O, Shekhar N, Gill CJ, Salami H, et al. Assessment of the Quality of Antenatal Care Services Provided by Health Workers Using a Mobile Phone Decision Support Application in Northern Nigeria: A Pre/Post-Intervention Study. *PLoS One*. 2015 May 5;10(5):e0123940.