

Ovarian Vein Thrombosis Post-Hysterectomy

A literature review of risk factors, incidence, implications, and management

Hallikeri. A, Vaselli. N, M, Walker. G

Introduction

Ovarian vein thrombosis (OVT) is rare type of venous thromboembolism, that occurs in 0.05-0.18% of vaginal births and in 2% of caesarean deliveries(1), however its prevalence post hysterectomy is still widely debated. This literature review aims to discuss risk factors, implications and management of this venous thromboembolic event.

Method

A literature review was conducted on PubMed and Google Scholar. The search terms ‘ovarian vein thrombosis’, ‘hysterectomy complications’ and ‘venous thrombosis’ were used to identify peer-reviewed and original articles published in English between 1990 – 2022. Articles were analysed and literature was categorised according to clinical presentation, investigation, prevalence, complications and management.




Image 1: Ovarian vein anatomy within the abdomen (6)
Illustration demonstrates ovarian veins arising from the ovarian venous plexus.



Image 2: CT scan of right ovarian vein thrombosis (7)
An abdominal-pelvic CT scan showing a hypodense tubular structure with peripheral enhancement to the right of the inferior vena cava, suggestive of thrombosis of the right ovarian vein.(1)

Pathophysiology

Anatomically, the ovarian veins originate from the venous plexus in the broad ligament and communicate with the uterine plexus. The ovarian vein on each side is in the retroperitoneum anterior to the psoas muscle (image 1). The right ovarian vein drains into the anterolateral IVC whilst the left vein drains into the left renal vein(2). The pathophysiology of OVT is ascribed to Virchows’ Triad of circulatory stasis, vessel wall injury and hypercoagulability(3). In 70-80% of cases, the right ovarian vein is implicated; this is likely due to its longer length and multiple incompetent valves(4). The biggest risk factors include; malignancy (44%), abdominal and pelvic surgery (33%), pelvic infection (17%) and pregnancy/puerperium (11%)(5).

Results

OVT can present as acute abdomen or an asymptomatic incidental finding upon imaging. Complications include pulmonary embolism, renal failure, sepsis and even some case reports of death(8). The gold standard investigations of choice differs between papers. The diagnosis of OVT can be established using ultrasound, MRI or CT. Ultrasound can be used as the first line of imaging as it is cost effective and without radiation risks, however, ovarian veins can be difficult to visualise in the presence of obesity and overlying bowel gas(9). Studies show MRI has the best combined sensitivity and specificity, as it is able to image in multiple planes and is sensitive to blood flow, it does not require IV contrast to reveal acute and subacute thrombi. MRI, however, is more expensive and time consuming than a CT. Dilated, thick-walled ovarian veins with rim enhancement and central hypodensity are considered to be the main CT imaging findings(10). Conversely, two papers state that laparotomy is the gold standard(11,12). Although multiple modalities can be used for diagnosis, contrast-enhanced CT studies seem to be the most cost effective and reliable technique(13). There remains debate over the gold standard of diagnosis and treatment of this condition. To date there are no official guidelines on the management of OVT. It can resolve spontaneously, however considering life-threatening complications, some studies encourage treatment with anti-coagulation(14). A recent study investigated the use of direct oral anticoagulants (DOACs) compared to warfarin and enoxaparin; they concluded that DOACs showed a similar risk-benefit ratio(5). The duration of anticoagulation has been suggested to be similar to that of lower extremity DVT; 3 months for provoked and 6 months for unprovoked(4,14,15,16). It is important to note that up to 50% of OVT cases may have a prothrombotic predisposition(17), therefore it is important that this is investigated and expert advice is sought as lifelong anticoagulation may be required. The use of antibiotics is recommended when there are complications such as ovarian abscess, thrombophlebitis, or sepsis(4,16). Surgical options are now less commonly used with the introduction of anticoagulation(16).

Conclusion

OVT is a serious condition with potential life-threatening complications. There remains debate over whether incidental/asymptomatic OVT should be treated unless there are complications. To date there remains no guidelines on the preferred modality of investigation or management of OVT. This literature review highlights the need for collaboration between different specialities to create a guideline on the optimal management of OVT for clinicians world-wide to follow.

References

1) Kodali N, Veytsman I, Martyr S, Lu K. Diagnosis and management of ovarian vein thrombosis in a healthy individual: a case report and a literature review. J Thromb Haemost [Internet]. 2017 Feb 1 [cited 2022 Jul 17];15(2):242–5. Available from: <https://pubmed.ncbi.nlm.nih.gov/27930855/>

2) Glick Y, Maingard J. Gonadal vein. In: Radiopaedia.org [Internet]. Radiopaedia.org; 2013 [cited 2022 Feb 20]. Available from: <http://radiopaedia.org/articles/26030>

3) O'Hagan S, Andronikou S, Truter R, Van Rensburg CJ. Ovarian vein thrombosis – a rare but important complication of hysterectomy and oophorectomy. S Afr J Obstet Gynaecol [Internet]. 2013 Sep 3 [cited 2022 Feb 20];19(3):86. Available from: <http://hmgp.co.za/index.php/saogp/article/view/1540>

4) Harris K, Mehta S, Ishakov E, Chalhoub M, Maniatis T, Forte F, et al. Ovarian vein thrombosis in the nonpregnant woman: an overlooked diagnosis. Ther Adv Hematol [Internet]. 2012 Oct [cited 2022 Feb 20];3(5):325–8. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/23616919>

5) Covut F, Kewan TZ, Perez O, Awada H, Babar A, Thapa B, et al. Direct oral anticoagulants (DOAC) versus warfarin and enoxaparin in ovarian vein thrombosis. J Clin Oncol [Internet]. 2019 May 20 [cited 2022 Feb 20];37(15_suppl):e14698–e14698. Available from: http://ascopubs.org/doi/10.1200/JCO.2019.37.15_suppl.e14698

6) El-Dasty M. Ovarian vein thrombosis: an overview for radiologist. In: ECR 2018 [Internet]. 2018 [cited 2023 May 28]. Available from: <https://epos.myscr.com/poster/esr/ecr2018/C-2058>

7) Martins MM, Carneiro C, Leibo S, Costa C. Ovarian vein thrombosis: a rare but not to be forgotten cause of abdominal pain. BMJ Case Rep [Internet]. 2021 Dec 30 [cited 2023 May 28];14(12):e246542. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/34969801>

8) Tanaka Y, Kato H, Hosoi A, Isobe M, Koyama S, Shiki Y. Ovarian vein thrombosis following total laparoscopic hysterectomy. Asian J Endosc Surg [Internet]. 2012 Nov [cited 2022 Feb 20];5(4):179–82. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/23095296>

References

9) Riva N, Calleja-Aguil J. Ovarian Vein Thrombosis: A Narrative Review. Hamostaseologie [Internet]. 2021 Aug [cited 2022 Feb 20];41(4):257–66. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/33348392>

10) Lebowitz JA, Rofsky NM, Krinsky GA, Weinreb JC. Gadolinium-enhanced body MR venography with subtraction technique. AJR Am J Roentgenol [Internet]. 1997 [cited 2022 Jul 17];169(3):755–8. Available from: <https://pubmed.ncbi.nlm.nih.gov/9275892/>

11) Dessole S, Capobianco G, Arru A, Demurtas P, Ambrosini G. Postpartum ovarian vein thrombosis: an unpredictable event: two case reports and review of the literature. Arch Gynecol Obstet [Internet]. 2003 [cited 2022 Jul 17];267(4):242–6. Available from: <https://pubmed.ncbi.nlm.nih.gov/12592429/>

12) Kominiares MA, Hibbard JU. Postpartum ovarian vein thrombosis: an update. Obstet Gynecol Surv [Internet]. 2006 May [cited 2022 Jul 17];61(5):337–42. Available from: <https://pubmed.ncbi.nlm.nih.gov/16635274/>

13) Kodali N, Veytsman I, Martyr S, Lu K. Diagnosis and management of ovarian vein thrombosis in a healthy individual: a case report and a literature review. J Thromb Haemost [Internet]. 2017 Feb 1 [cited 2022 Jul 17];15(2):242–5. Available from: <https://pubmed.ncbi.nlm.nih.gov/27930855/>

14) Trang N, Kalluri M, Bajaj T, Petersen G. Idiopathic Left Ovarian Vein Thrombosis. J Investig Med high impact case reports [Internet]. 2020 Jan 1 [cited 2022 Feb 20];8:2324709620947257. Available from: <http://journals.sagepub.com/doi/10.1177/2324709620947257>

15) Wysokinska E, Hodge D, McBane R. Ovarian vein thrombosis: incidence of recurrent venous thromboembolism and survival. Thromb Haemost. 2006;Aug;96(2):126–31.

16) Plastini T, Henry D, Dunleavy K. Ovarian vein thrombus: to treat or not to treat? Blood Adv [Internet]. 2017 Jun 27 [cited 2022 Feb 20];1(15):1120–3. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/29296754>

17) Salomon O, Apter S, Shaham D, Hiller N, Bar-Ziv J, Itzchak Y, Gitel S, Rosenberg N, Strauss S, Kaufman N SU. Risk factors associated with postpartum ovarian vein thrombosis. Thromb Haemost. 1990;Sep;82(3):1015–9.