

Compliance and prediction of same day discharge after laparoscopic hysterectomy: A retrospective cohort study

BACKGROUND

A standardised same day discharge pathway for laparoscopic hysterectomy was introduced at the Birmingham Women’s and Children’s Hospital in 2019. Our pre-, intra- and post-operative protocols have been refined since then with the aim of enhancing successful same day discharge.

OBJECTIVE


- To assess rates of compliance with a same day discharge pathway.
- To confirm whether previously reported factors predictive of successful same day discharge following laparoscopic hysterectomy were valid in at Birmingham Women’s Hospital, a UK teaching hospital.

METHODS AND DESIGN

- Retrospective cohort study of 186 patients undergoing laparoscopic hysterectomy between January 2022-December 2024 (24 months) who were planned for same day discharge
- Eligible patients were identified using Operating Management Information System (ORMIS)
- The primary outcome was the rates of successful same day discharge
- Multivariable logistic regression analysis was used to assess factors considered potentially prognostic for successful same day discharge. Independent variables include:
 - Patient characteristics - age, ethnicity BMI, ASA status, previous abdominal surgery, smoking status and distance to home.
 - Surgical variables - blood loss, concomitant surgery, duration, adhesiolysis, operation time and pneumoperitoneum pressure.

RESULTS

- 132/186 (70.9%) women were discharged from the hospital on the same day.
- Two factors were found to be significantly predictive of successful same day discharge:**
 - Timing of surgery before 14:00:** 73% (122/168) before 14:00 vs. 25% (2/8) from 14:00 onwards; p=0.01)
 - Pneumoperitoneum >/= 10mmHg:** 73% >/=10mmHg (120/165) vs. 56% <10mmHg; (10/18) (p=0.01)



DISCUSSION

- Patients suitable for same day discharge should be scheduled for surgery before 14:00
- Low pressure surgery, using operating pressures <10mmHg, has been advocated to reduce post-operative pain. However, in our series low insufflation pressures reduced the chance of same day discharge.
- Larger, multicentre datasets are needed to confirm these findings, explore optimal operating pressures and evaluate post-operative factors.
- Variables should also be evaluated that predict rates of readmission.