

Incidence of maternal and fetal/neonatal outcomes in morbidly obese parturients undergoing spinal anesthesia (SAB) for elective cesarean delivery (CD) at term

Tianna Xia, DO; Ling-Qun Hu, MD; Jason Hoang, BS; Brett Worly, MD; Blair Hayes, MD; Goran Ristev, MD; Yun Xia, MD, PhD

INTRODUCTION

- Most CD are performed under neuraxial anesthesia in LUD
- BP measurements taken on the arm may not reflect uterine perfusion pressure
- Risk factors for poor fetal/neonatal outcomes in otherwise healthy fetuses may include maternal obesity and HTN^[1] but need further investigation
- It is valuable to the field of OB anesthesia to determine the relationship between incidence of poor maternal and fetal/neonatal outcomes and average BP after SAB

METHODS

- **Design:** Single-center, prospective, cohort study
- **Population:** Age ≥ 18 , BMI ≥ 35 , ASA II-III, elective CD
- **Aim:** Characterizing BP at the left ankle instead of upper arm
- **Null hypothesis:** BP measurements taken at the left ankle = BP measurements taken on the arm
- **Primary outcome:** Incidence of maternal hypotension
- **Secondary outcomes:**
 - Maternal: arm comfort, nausea/vomiting, shivering
 - Neonatal: FHR, APGAR scores, airway support, NICU admission

Table 1. Incidence of maternal and fetal/neonatal outcomes

Primary Outcomes (n (%))			
● Hypotension after SAB at non-IV arm		● Hypotension after SAB at left ankle	
➤ In supine	15 (31%)	➤ In supine	6 (12%)
➤ In LUD	18 (37%)	➤ In LUD	6 (12%)
➤ In supine after newborn	8 (16%)	➤ In supine after newborn	2 (4%)
➤ In LUD after newborn	8 (16%)	➤ In LUD after newborn	4 (8%)
Secondary Outcomes			
● Maternal:		● Fetal/Neonatal (n (%)):	
○ Nausea/vomiting (n (%))	9 (18%)	○ FHR ≤ 110 after spinal	5 (10%)
○ Shivering (n (%))	9 (18%)	○ Airway support (facemask, CPAP, O2, intubation)	2 (4%)
○ Pain Arm > Ankle during measuring (n (%))	6 (12%)	○ NICU admission	7 (14%)
○ Mean SBP before SAB (mmHg, mean (SD))		○ Apgar Score at 1 min	
➤ Arm in supine	136 (19)	≤ 3	5 (10%)
➤ Arm in LUD	133 (18)	> 3 ~≤ 7	5 (10%)
➤ Left ankle in supine	153 (20)	> 7	39 (80%)
➤ Left ankle in LUD	159 (18)		
○ Mean SBP after SAB (mmHg, mean (SD))		○ Apgar Score at 5 min	
➤ Arm in supine	122 (20)	≤ 3	0 (0%)
➤ Arm in LUD	118 (22)	> 3 ~≤ 7	3 (6%)
➤ Left ankle in supine	143 (28)	> 7	46 (94%)
➤ Left ankle in LUD	141 (27)		
○ Mean SBP after delivery (mmHg, mean (SD))		○ Composite fetal/neonatal outcome	14 (29%)
➤ Arm in supine	121 (13)	(either FHR ≤ 110, or Apgar ≤ 7 at 1 min, or 5 min, or airway support, or NICU admission)	
➤ Arm in LUD	120(13)		
➤ Left ankle in supine	139 (22)		
➤ Left ankle in LUD	138 (23)		

RESULTS

➤ **Demographics**

- 32±5 yo
- 38w3d±1.3 GA
- 82% ASA III
- BMI 44±7.8
- 31% HTN

➤ **Maternal hypotension after spinal**

- LUD at arm: 37%
- Supine at arm: 31%
- LUD/supine at left ankle: 12%

➤ **Fetal & neonatal outcomes**

- Composite: 29%
- NICU admission rate: 14%

➤ LUD by tilting OR table 15 degrees is not realistic

FUTURE STEPS

- Investigate the relationship between poor neonatal outcomes and the current definition of maternal hypotension
- Correlate hypotensive BP readings with poor outcomes and identify the best BP measurement site for predicting prognoses
- Address relationship of timing of position changes as it relates to BP
- Evaluate the association between time from SAB to delivery and poor neonatal outcomes

REFERENCES

1. Obes Res Clin Pract. Nov-Dec 2021;15(6):536-545
2. Acta Anaesthesiol Scand 2010: 54, 8, 909-921
3. ACOG 2008:111,4, 823-828