

Reduction of General Anesthesia Rate by Improving Multidisciplinary Communication Before Unscheduled Cesarean Sections



Andrea Girnius MD¹, Candice Snyder MD², Heather Czarny MD², Tom Minges CRNA¹, Michael Stacey CRNA¹, Tamara Supinski RN³, Muhammad A Zafar, MD, MSc⁴



¹Department of Anesthesiology, University of Cincinnati Medical Center; ²Department of Obstetrics and Gynecology, University of Cincinnati Medical Center; ³ Labor and Delivery Nurse Manager, University of Cincinnati Medical Center; ⁴Department of Pulmonary and Critical Care Medicine, University of Cincinnati Medical Center

Introduction:

- The use of general anesthesia (GA) for cesarean section (C/S) is associated with elevated maternal risk¹.
- Our overall GA rate for C/S ranged from 9-17%, with GA use for unscheduled C/S specifically as high as 22% in 2020.
- Failures of communication in healthcare are common and can lead to suboptimal patient care as well as staff dissatisfaction and poor work environment^{2,3}.
- On labor and delivery, communication about unscheduled C/S was often fragmented and incomplete, which can contribute to avoidable use of GA.

Methods:

- Using the Model for Improvement, we sought to understand the failures leading up to unscheduled C/S that may lead to avoidable GA use.
- We found that communication among nursing, anesthesia, and obstetric (OB) teams was fragmented, delayed, and frequently incomplete and inadequate for providing coordinated multidisciplinary care.
- We developed and introduced a **multidisciplinary, patient centered pre-C/S huddle** attended by representatives from nursing, anesthesia, and OB.
- A standard checklist of critical information was used as a discussion guide. It was tested and refined through iterative PDSA cycles prior to implementation.




Fig 2: Pareto Chart

Pre-C Section Huddle: UNSCHEDULED

Location: Patient's room

Attendees:

- ☐ OB R4/R3 resident
- ☐ Patient's RN
- ☐ Charge RN
- ☐ Anesthesia representative

Time spent in huddle:

Discussion Checklist:

- ☐ Indication for cesarean
- ☐ Decision time
- ☐ Classification of cesarean- unscheduled vs emergent
- ☐ Agreement on expected timing: Define goal time in the OR
- ☐ Mode of anesthesia
- ☐ Antibiotics requested/who will obtain them
- ☐ Peds needs (NICU, Fetal Care) or other key neonatal information
- ☐ Which MDs are performing the cesarean section
- ☐ Acknowledgement that everyone in agreement with plan
 - If there are concerns for safety that are not appropriate to discuss in front of the patient, then the phrase, "LET'S STEP OUT TO DISCUSS LOGISTICS" can be used to alert the team that a discussion outside of the room needs to occur.

Fig 3: Huddle Checklist

Measures:

Outcome measures:

- GA rate for unscheduled C/S
- Satisfaction with communication regarding C/S plan among anesthesia, nursing, and OB teams.

Process measures:

- Adherence to pre-C/S multidisciplinary team huddle.

Balancing measure:

- Decision to incision interval (DTI) for unscheduled C/S.

	Process 1	Process 2	Process 3
Changes Tested	Patient Centered pre C/S huddles to improve communication	Data Collection/Huddle completion tracking	Increase Staff Engagement in huddle process
# of PDSA's/ Ramps	2 Ramps 11 PDSA's	2 Ramps 12 PDSA's	3 Ramps 11 PDSA's
Results	Huddles improved communication between teams, reduced GA for unscheduled C/S, and were well received by patients	Successful measures included anesthesia team returning huddle checklist and noting # of C/S and huddles at shift change.	Most measures increased knowledge/awareness of huddles. Review at mid-shift safety huddle facilitated discussion of missed opportunities

Table 1: Summary of Plan-Do-Study-Act (PDSA) cycles

Conclusions:

- System barriers led to unclear and inaccurate communication surrounding unscheduled C/S.
- Implementation of a **multidisciplinary in-person pre C/S huddle** improved communication between teams and reduced general anesthesia rates for unscheduled C/S.
- Our next steps are to ensure sustainability of the process and continue to mitigate other causes of avoidable GA use for C/S.

References:

- Mushambi et al. *Anaesthesia*. 2015;70(11):1286-1306.
- Lingard LS et al. *Qual Saf Health Care*. 2004;13:330-334
- Shortell SM et al. *Med Care*. 1994;32:508-525

Results:

