


Labor Epidural Anesthesia Neuraxial Outcomes in Super Morbidly Obese Patients as compared to General Population



Ahmed Butt, Mauricio Ramos Lozano, Safir Abbas, Trent Weatherley, Kristine Lane, Haley Hodgins, Patrick Finnegan, Hiram Acevedo, MD, Michele Simon, MD, Rovnat Babazade, MD
Department of Anesthesiology
University of Texas Medical Branch, Galveston, TX



Introduction

When placed successfully, epidural anesthesia does not carry significant risk in cesarean sections and immediate newborn health (1). While neuraxial anesthesia is often difficult to place in obese patients, it allows the patient to avoid general anesthesia which reduces airway manipulation, curtails the risk of aspiration, and decreases volatile anesthetic exposure to both the fetus and the parturient. (2)

The present study aims to present and analyze the neuraxial outcomes in the super obese patients who received labor epidural anesthesia.

Materials & Methods (Operative Course)

We conducted a retrospective review of patients who presented with BMI $\geq$ 50.0 upon admission for delivery at a single hospital institution. We collected data from the electronic medical records at to identify patients who were admitted for delivery of a viable fetus between 2015-2021. We reviewed 342 patient encounters, of which 108 received epidural analgesia (32% of the observed population). Data collected included the parturients' demographic information and their anesthesia information for each L&D admission.

Results

Table 1 details the demographics of the patient population and their anesthetic outcomes. Of the 108 patients, maternal age averaged at 28.33 with a SD of 5.97 and maternal BMI averaged at 55.47 with an SD of 5.92. Of the 108 epidurals replaced, 0.90% (1/108) required faculty intervention, 11.65% (12/108) required replacement, 3.70% (4/108) resulted in failure, 2.78% (3/108) were suspected of having dural puncture and 3.70% (4/82) reported post dural punctual headaches. Overall, 24.07% (26/108) of epidurals had complications.

Table 1. Super Obese Parturients Who Received Epidural Anesthesia Outcomes			
Subjects Included (n)			108
Maternal Age (years)	Mean		28.33
	SD		5.97
	Minimum		17
	Maximum		42
Maternal BMI (kg/m ²)	Mean		55.47
	SD		5.92
	Minimum		50.1
	Maximum		92.23
Neuraxial Outcomes (n) (of 81 charts)	US Guidance Used		2
	Required Faculty Intervention		1
	Neuraxial Failure Requiring Anesthesia Mode Change (ie to Spinal, CSE, or GETA)		4
	Required Epidural Replacement		12
	Suspected Dural Puncture		3
	Post Dural Puncture Headache		4

Discussion

Although some of the epidurals placed in our obese population resulted in complications, most were placed successfully without complications. Reducing the number of epidural complications can improve a patient's labor experience by reducing the number of headaches experienced, minimize the use of extra equipment that has to be used in neuraxial replacement, and prevent use of general anesthesia and volatile anesthetics in C-sections.

Discussion (cont.)

Moving forward, we would need to obtain a control group that is truly representative of the general population to see if the number of complicated cases that we obtained is statistically significant in comparison. If our sample showed statistical significance, the next step would involve investigation of the effect of additional interventions, such as ultrasound and other assist devices, on the reduction of epidural complications.

References

- Anim-Somuah M, Smyth RM, Cyna AM, Cuthbert A. Epidural versus non-epidural or no analgesia for pain management in labour. *Cochrane Database Syst Rev*. 2018;5(5):CD000331. Published 2018 May 21.
doi:10.1002/14651858.CD000331.pub4
Taylor CR, Dominguez JE, Habib AS. Obesity And Obstetric Anesthesia: Current Insights. *Local Reg Anesth*. 2019;12:111-124. Published 2019 Nov 18. doi:10.2147/LRA.S186530