

Cardiovascular Collapse, Multi-Chamber Intracardiac Thrombosis, and Disseminated Intravascular Coagulation Following Dilation & Evacuation in a Post-Partum Cardiomyopathy Patient

Hilary Gallin, MD, MBA, Rebecca Minehart, MD, and Marvin Chang, MD
Department of Anesthesia, Critical Care, and Pain Medicine
Massachusetts General Hospital, Boston, Massachusetts, USA

Peripartum cardiomyopathy (PPCM) is associated with 11% of maternal deaths,¹ both of which can be mitigated by performing a D&E in the highest risk patients. D&E are low risk procedures often performed in the office setting with low complication rates.

Patient History

- ▶ 32yo female G4P3 at 16 weeks with past medical history notable for asthma and prior asymptomatic COVID was transferred from OSH where she presented with 4 months of worsening shortness of breath.
- ▶ TTE revealed an LVEF 10%, moderately reduced RV function, moderate to severe MR, and RSVP of 60.
- ▶ Imaging was negative for PE, CAD, and DVT. She agreed to undergo a D&E to reverse her PPCM.
- ▶ Of note, she had tolerated a laparoscopic cholecystectomy under general anesthesia 3 weeks earlier.



¹Lewey J and Jennifer H. "Cardiomyopathy in pregnancy." *Seminars in perinatology* vol. 38.5 (2014): 309-17. 2014.04.021



Perioperative Course

Pre-Operative

- Hemodynamics, volume status, and rhythm control of ventricular tachyarrhythmias were optimized by Cardiology, and her prophylactic enoxaparin held the day before her D&E.
- Planned anesthetic for the D&E was MAC with a propofol infusion and ketamine.

Intraoperative

- Initial vitals were BP 117/79, HR 115, and SpO₂ 100%, and were stable during the case.
- Immediately upon D&E completion, SpO₂ fell to the 70's despite no change in her spontaneous breathing.

Emergency Response

- Positive pressure ventilation was started and chest compressions were initiated.
- She was placed on peripheral ECMO.
- Differential diagnosis at the time included
 - Amniotic fluid embolism
 - Pulmonary embolism
 - Venous air embolism
- TEE was notable for a PFO and diffuse clots in all chambers of her heart including at the interface of the IVC and RA.
- Labs were notable for an INR 2.5, platelets 52, and Fibrinogen 177.



Patient Outcome and Discussion

Postoperative Care

- She was transferred to the ICU.
- She underwent surgical thrombectomy, biventricular assist device (BiVAD), and central ECMO cannulation.
- Her neurologic status deteriorated with imaging revealing significant embolic infarcts and was deemed not a candidate for heart transplant.
- Care was withdrawn by the family on POD 14.

Discussion

- Our patient with cardiovascular collapse, multi-chamber intracardiac thrombosis, and DIC likely suffered an AFE, a rare complication of second trimester D&E.²
- Pregnant patients placed on extracorporeal life support (ECLS) have high survivability rates compared to other etiologies if the cause is primarily cardiac in nature.³
- In our patient, an embolic event may not have been preventable and likely decreased her survivability despite ECLS.³