

Anesthetic Planning for Delivery for a Patient with Titin-Truncating Variant Mutation Complicated by Complex Arrhythmia

Kiana C. Nguyen BS, Elliott Callahan MD, Peter Yeh MD

INTRODUCTION

- ◉ **Titin (TTN)**
 - Sarcomere organization
 - Striated muscle function
- ◉ **TTN-truncating variants (TTNtv)**
 - Genetic dilated cardiomyopathy¹
 - Peripartum cardiomyopathy¹
 - Early-onset arrhythmia²




Figure 1. 30w TTE, apical 4 chamber view. End-diastole (above) and end-systole (below). Pacer/ICD lead indicated with arrow.

CASE

- ◉ **27 yo G3P0 at 33w3d**
 - TTNtv c.87852G>A (p.Trp29284*)
- ◉ **PMH:**
 - Sudden cardiac arrest, age 16
 - Atrial fibrillation
 - AICD placed
- ◉ **MEDS:** quinidine, cilostazol, metoprolol
- ◉ **mWHO class II-III, CARPREG II score 5**

Anesthetic Planning for Delivery for a Patient with Titin-Truncating Variant Mutation Complicated by Complex Arrhythmia

Kiana C. Nguyen BS, Elliott Callahan MD, Peter Yeh MD

DELIVERY PLAN



Discontinue
cilostazol 48 hrs
prior to neuraxial
anesthesia



Manage potential
breakthrough
arrhythmia with
isoproterenol

Monitor in ICU while
holding cilostazol



IOL at 39 weeks
with neuraxial
anesthesia



LABOR COURSE

- ◉ Patient p/w SROM at 38w4d
- ◉ Insufficient window off cilostazol
 - Thus, no neuraxial anesthesia
- ◉ Pt received remifentanyl with pudendal block for 2nd stage analgesia
- ◉ Patient had an uncomplicated delivery
 - No episodes of arrhythmia


Anesthetic Planning for Delivery for a Patient with Titin-Truncating Variant Mutation Complicated by Complex Arrhythmia

Kiana C. Nguyen BS, Elliott Callahan MD, Peter Yeh MD

DISCUSSION

◉ Cilostazol

- PDE-3 inhibitor
- Anti-arrhythmic
- Inhibits platelet aggregation
- Contraindication to neuraxial procedures



◉ Early epidural in cardiac disease

- Avoid sympathetic surges associated with uterine contractions
- May help prevent precipitating malignant arrhythmia during labor

CONCLUSION

◉ Pregnancy is a cardiac stressor!

- “Second hit” in TTNtv patients at risk for PPCM & malignant arrhythmia¹

◉ Risk stratification

- Cardiac MRI assessing for midwall fibrosis is a known predictor of arrhythmic events and need for device therapy in DCM.³
- Explore future roles for cardiac MRI in PPCM

REFERENCES

1. Goli R, Li J, Brandimarto J, et al. Genetic and Phenotypic Landscape of Peripartum Cardiomyopathy. *Circulation*. 2021;143(19):1852-1862.
2. Choi SH, Weng LC, Roselli C, et al. Association Between Titin Loss-of-Function Variants and Early-Onset Atrial Fibrillation. *JAMA*. 2018;320(22):2354-2364.
3. Corden B, Jarman J, Whiffin N, et al. Association of Titin-Truncating Genetic Variants With Life-threatening Cardiac Arrhythmias in Patients With Dilated Cardiomyopathy and Implanted Defibrillators. *JAMA Netw Open*. 2019;2(6):e196520.