

Syrinx and Symptomatic Covid-19 in Pregnancy

Robert P Schroell III MD,¹ Leslie J Matthews MD, PharmD,¹ Blair H Hayes MD,¹
¹ Department of Anesthesiology, The Ohio State University, Wexner Medical Center, Columbus, Ohio, USA

Introduction

- Syringomyelia
 - A rare spinal cord pathology results in damage to the tissue
- Syrinx: fluid-filled cyst within the spinal cord
- Causes
 - Congenital
 - Traumatic
- Currently no consensus for management


Case Presentation

- 31yo F G3P0111 at 34w2d with history of scoliosis, traumatic MVC complicated by temporary paraplegia, syringomyelia with sensory deficits below T8-9 status post spinal fusion. History of 1 prior Cesarean delivery
- Presented with headaches and severe range blood pressures, diagnosed with pre-eclampsia with severe features
- Exam findings: shortness of breath, fatigue, congestion, frequent cough -- found to be COVID-19 positive

Imaging (MRI from 9 months prior)

- Syringomyelia with 2 discrete syrinx:
 1. C7 to T1
 2. T 2-3 to the filum terminale

Syrinx



Case Presentation: Imaging



Delivery Course

Decision Making

- Neurosurgery consult on admission
- Multidisciplinary team discussion
 - No time for additional imaging given urgency of delivery due to pre-eclampsia
- Uncomplicated CS under GA, extubated in OR
- Post op course uneventful, discharged home

Discussion

- Neuraxial anesthesia with syrinx can be risky
 - Changes seen in CSF flow dynamics
 - May cause additional syrinx
 - Devastating neurological consequences such as herniation
- Research has been limited to case studies
 - Most undergo GA
 - Recent imaging imperative
- COVID-19 in pregnancy
 - Advantages of neuraxial anesthesia
 - Provides excellent postoperative analgesia
 - Allows for early maternal/infant bonding
 - Maintains spontaneous ventilation, protected airway
 - Known difficult airway in pregnancy
 - Increased aspiration risk
- Concerns with airway manipulation in COVID-19
 - Increased risk of viral transmission
 - Positive Pressure ventilation in COVID-19 patients

References